



CG


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : G06F 19/00		(11) International Publication Number: WO 00/70528
A2		(43) International Publication Date: 23 November 2000 (23.11.00)
(21) International Application Number: PCT/US00/13154		(74) Agent: LOUIE, Michael, L.; Beyer Weaver & Thomas, LLP, P.O. Box 130, Mountain View, CA 94042-0130 (US).
(22) International Filing Date: 12 May 2000 (12.05.00)		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(30) Priority Data: 09/310,879 14 May 1999 (14.05.99) US 09/311,890 14 May 1999 (14.05.99) US 09/311,996 14 May 1999 (14.05.99) US 60/134,104 14 May 1999 (14.05.99) US Not furnished 26 April 2000 (26.04.00) US		
(71) Applicant: CYTOKINETICS, INC. [US/US]; Suite 2, 280 East Grand Avenue, South San Francisco, CA 94080 (US).		
(72) Inventors: SABRY, James, H.; 52 Buena Vista Terrace, San Francisco, CA 94117 (US). ADAMS, Cynthia, L.; 615 Georgia Avenue, Palo Alto, CA 94306 (US). VAISBERG, Eugeni, A.; 647 Pegasus Lane, Foster City, CA 94404 (US). CROMPTON, Anne, M.; 2 Bellair Place, San Francisco, CA 94133 (US). BLUM, Robert, I.; 17 Shoreview Avenue, San Francisco, CA 94121 (US). OESTREICHER, Donald, R.; 904 Old Town Court, Cupertino, CA 95014-4024 (US). SIGAL, Nolan, H.; 941 Berry Avenue, Los Altos, CA 94024 (US).		
		Published Without international search report and to be republished upon receipt of that report.

(54) Title: METHOD AND APPARATUS FOR PREDICTIVE CELLULAR BIOINFORMATICS

(57) Abstract

Techniques for using information technology in therapeutics or drug discovery. In an exemplary embodiment, techniques for determining information about the properties of substances based upon information about structure of living or non-living cells exposed to substances are provided. A method according to the present invention enables researchers and/or scientists to identify promising candidates in the search for new and better medicines or treatments using, for example, a cellular informatics database. The present invention further teaches a system for acquiring knowledge from cellular information. The system has a database 1012 comprising a database management module ("DBMS"). The system also has a variety of modules, including a population module coupled to the DBMS for categorizing and storing a plurality of features (e.g., cell size, distance between cells, cell population, cell type) from an image acquisition device into the database. The system has a translation module coupled to the DBMS for defining a descriptor from a set of selected features from the plurality of features. In a specific embodiment, the descriptor is for a known or unknown compound, e.g., drug. A prediction module is coupled to the DBMS for selecting one of a plurality of descriptors from known and unknown compounds from the database based upon a selected descriptor from a selected compound. The selected compound may be one that is useful for treatment of human beings or the like.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

PATENT APPLICATION
METHOD AND APPARATUS FOR
PREDICTIVE CELLULAR BIOINFORMATICS
COPYRIGHT NOTICE

5 A portion of the disclosure of this patent document contains material which is subject to copyright protection. The copyright owner has no objection to the facsimile reproduction by anyone of the patent document or the patent disclosure as it appears in the Patent and Trademark Office patent file or records, but otherwise reserves all copyright rights whatsoever. The present description includes examples of
10 computer codes, which may be used to implement aspects of the present invention. Assignee of the present invention reserves all rights with respect to these codes and provides notice herein. Notice is hereby given © Cytokinetics, Inc. 1999.

BACKGROUND OF THE INVENTION

 The present invention provides techniques for information
15 management using a database platform. More particularly, the present invention provides a system including computer code that couples to a database device. The system provides for image capturing of living, dead, or fixed cells or cell fractions used to identify information about substances used on the cells or information about the cells themselves. Accordingly, the present invention can enable researchers and
20 scientists to identify promising candidates in the search for new and better medicines, for example, in drug discovery and development. The principles enumerated herein may, with equal facility, be applied to other applications, including but not limited to use in environmental applications such as determining chemical toxicities and other non-pharmaceutical toxicology uses.

25 For a long time, researchers in the pharmaceutical field have sought for better ways of searching for substances possessing properties that make them suitable as medicines. In the early days, researchers generally relied upon extracts from plants, dyes, and microbiological extracts for such substances. Examples of such substances include the pain reliever aspirin, the anti-cancer drug paclitaxel (brand
30 name TaxolTM), and the heart medication called digoxin. The number of useful medicines has generally been limited.

Purified substances having desirable bio-active properties are also often difficult to discover. Advances in traditional organic chemistry and more recently the rapid chemical synthesis methods often referred to as combinatorial chemistry have increased the number of compounds that researchers test for biological activity. Originally, substances were often initially tested on animals or humans to determine their biological activity. While results from such tests may identify a good drug candidate, they are often time consuming and costly, thus a limited number of substances can be tested. Therefore, pharmaceutical companies have turned to testing their ever-increasing libraries of substances against isolated proteins (drug targets) in biochemical assays that can be carried out at high throughput and low cost. It should be noted that the substances need to be tested in numerous protein tests, each customized for a particular drug target. Therefore, although each protein test may be run at a high-throughput, the design of multiple protein tests can be time-consuming. Substances deemed promising based on results from the protein tests are then tested in lower throughput cellular and animal tests.

There have been some attempts to use image acquisition techniques to screen a large number of substances based upon biological cell information. One such attempt is described in International Application No. WO 98/38490 in the names of Dunlay, et al. Dunlay et al. generally describes a conventional image acquisition system. This conventional system collects and saves images based on certain criteria that are predefined, not on a fixed area of an imaging surface. Additionally, the conventional system has poor lighting design, which makes image processing for multiple cells difficult. Furthermore, the conventional system is not designed for capturing, populating and utilizing a large database design. The conventional system is designed for customized cellular assays, not as a tool for generation of a cellular informatics database. Without such database capabilities the conventional system cannot be used for screening, analyzing, and comparing large quantities of cells from multiple experiments on multiple days in a predictive, efficient and cost effective manner.

What is needed is a rapid assay to assess the activity of compounds against multiple drug targets simultaneously in a cellular context. What is also needed are techniques for finding the effects of substances on cell function based upon searching and analyzing cellular information.

SUMMARY OF THE INVENTION

According to at least one embodiment of the present invention, techniques for determining information about effects of potential substances on cells are provided. In another exemplary embodiment, the present invention provides a novel system including hardware, computer codes, user interfaces, and a database for acquiring, storing and retrieving cellular and substance information. The cells can include living, dead, or fixed cells or fractions of cells. The present invention enables, *inter alia*, researchers and/or scientists to identify promising candidates in the search for new and better medicines or treatments using, for example, a cellular informatics database.

According to the present invention, a computer program for identification and verification of biological properties of substances can include code that causes a sample of a substance to be administered to a cell. The code determines one or more features for two or more cell components, or markers, in the presence of the substance. The code can form one or more descriptors from the features. Descriptors can be formed by combining features of two or more cell components as identified using the markers. The code can then search one or more descriptors obtained from prior administered substances upon cells in order to locate descriptors having a relationship to the descriptors noted for the substance under study. The code predicts properties of the administered substance based upon the properties of the prior administered substances using the relationship between the descriptors. The code can provide for identifying properties of substances based upon effects on cell characteristics. Candidate drug mechanisms of action, potency, specificity, pharmacodynamic, and pharmacokinetic parameters, toxicity, and the like can be used as substance properties.

In a specific embodiment, the present invention provides a system for acquiring knowledge from cellular information. The system has a database comprising a database management module ("DBMS"). The system also has a variety of other modules, including a population module that is coupled to the DBMS and serves to categorize and store a plurality of features (including but not limited to cell size, distance between cells, cell population, as well as sub-cellular features such as organelle location, protein location and sub-cellular constituent location and

movement) from an image acquisition device into the database. The system has a translation module coupled to the DBMS for defining a descriptor from a set of selected features from the plurality of features. In a specific embodiment, the descriptor is for a known or unknown compound, e.g., drug. A prediction module is
5 coupled to the DBMS for selecting one of a plurality of a descriptors from known and unknown compounds from the database based upon a selected descriptor from a selected compound. The selected compound may be one that is useful for treatment of human beings or the like.

In a specific embodiment, the present invention provides a system for
10 populating a database with cellular information. The system includes a cell holder (e.g., multi-well plate, chip, microfluidic assembly, or other cell chamber) comprising a plurality of sites in a spatial orientation. Each of the sites is capable of holding a plurality of cells to be imaged. Note – the light guide is one embodiment, but we don't want to be limited to it.

15 According to one embodiment, the present system also has an illumination apparatus including a liquid light guide operably coupled to the imaging device for highlighting the plurality of cells in a relatively even spatial manner for image capturing and measurement purposes. Still further, the liquid light guide allows sub-elements (e.g., filter, lamp) of the illumination apparatus to be placed at a
20 remote location to prevent mechanical interference of the cell holder during image capturing. Alternative lighting methodologies may, with equal facility, be implemented.

The system also has an image-capturing device (e.g., charge coupled device camera, translation stage, shutter, microscope, software, shutter control) coupled to a
25 computing device (e.g., computer, network computer, work station, analog computing device, on-board image-processor, and laptop). The image-capturing device is adapted to capture at least one image in at least one of the plurality of sites. One some embodiments, multiple images can be captured, where each image represents a different cell component (or portion). The image-capturing device can be adapted to
30 convert the image into a digital representation, which highlights the feature or features of the one site.

A database storage device (e.g., relational database, object oriented database, mixed object oriented database) includes a database management element. The

database is coupled to the image capturing device. In a specific embodiment, the present system includes modules for feature extraction, generation of descriptions, and data preparation and analysis.

In a specific embodiment, the present invention provides a novel
5 system for determining an effect of a manipulation of a cell using one or more image frames. The system has a plate comprising a plurality of sites in a spatial orientation. Each of the sites is capable of holding a plurality of cells to be imaged. The system also has an image capturing device to capture a plurality of images of at least one site from the plurality of sites. The image capturing device is coupled to the computing
10 device. The system also has an image processing device to combine the plurality of images of at least one site or plurality of sites. The image processing device is operably coupled to the plate. An image processing device is also included. The image processing device can be adapted to form a digitized representation of the plurality of images from the site or plurality of sites. Furthermore, the system has a
15 database storage device comprising a database management element. The database can be adapted to retrieve the descriptor or descriptors of the plurality of features from the computing processing device and storing them in a selected manner.

In a specific embodiment, the present invention provides a system for capturing cellular information. The system also has an image acquisition system
20 comprising a charged coupled device camera adapted to capture an image of a plurality of manipulated cells in various stages of the cell cycle. The stages of the cell cycle are currently understood to include interphase, G0 phase, G1 phase, S phase, G2 phase, M phase, prophase, prometaphase, metaphase, anaphase, and telophase. The principles of the present invention specifically contemplate the application thereof on
25 additional cell cycle stages when and if they are identified.

An optical source is coupled to the image acquisition system for highlighting the plurality of manipulated cells in the various stages of the cell cycle. The illumination apparatus provides for an acquisition of the image of the plurality of manipulated cells. In a specific embodiment, the illumination apparatus has a liquid
30 light guide coupled to a light source at a remote location.

A variety of user interfaces are utile for accessing the several features of the present invention. Those having ordinary skill in the art will appreciate that different user interfaces may be required to support different research scenarios. The

present invention specifically contemplates the utilization of a wide variety of user interfaces.

Numerous benefits are achieved by way of the present invention over conventional techniques. The present invention can provide techniques for predictive
5 cellular bioinformatics that can streamline a number of important decisions made in the drug discovery industry. The present invention can be implemented using off the shelf hardware including databases. In other aspects, the present invention can find useful information about substances as well as cells or portions of cells. Furthermore, the present invention can acquire more than one feature using more than one
10 manipulation. Moreover, the present invention can provide information about a wide variety of cellular information that is not conventionally available. This information includes information about different cell components, e.g., nuclei and Golgi apparatus. Still further, the present invention provides an automated or semi-automated technique for acquiring images and populating a database. The present
15 database can be combined with others such as genomics, and the like. Moreover, the present invention can be implemented to predict, *inter alia*, a mechanism of action, toxicity, target validation, and pre-clinical disease model.

A further understanding of the nature and advantages of the invention herein may be realized by reference to the remaining sections of the specification and
20 the attached drawings.

BRIEF DESCRIPTION OF THE DRAWING

For more complete understanding of the present invention, reference is
5 made to the accompanying Drawing in the following Detailed Description of the
Invention. In the drawing:

Fig. 1 is a simplified system diagram according to an embodiment
according to the present invention;

Figs. 1A-1B are more detailed diagrams of database systems according
10 to embodiments of the present invention;

Fig. 2 is a simplified block diagram according to an alternative
embodiment according to the present invention;

Figs. 3-6 are simplified diagrams of system elements according to
embodiments of the present invention;

15 Figs. 7A-7K illustrate representative block diagrams of simplified
process steps in a particular embodiment according to the present invention;

Fig. 8A-8F illustrate representative quantified descriptors of effects of
manipulations on images of cells in a particular experiment;

Fig. 9 illustrates example images for different types of morphologies in
20 a particular experiment;

Fig. 10 illustrates a distribution of various morphologies in a cell
population responsive to drug concentration in a particular experiment;

Fig. 11 illustrates a graph of quantified features of effects of
manipulations on cells in a particular experiment;

25 Fig. 12 illustrates effects of external agents on cells in a particular
experiment;

Fig. 13 illustrates 4 panels for each marker for a plurality of A549 cells
in a particular experiment;

Fig. 14 illustrates 4 panels for each marker for a plurality of OVCAR-3
30 cells in a particular experiment;

Fig. 15 illustrates 4 panels for each marker for a plurality of OVCAR-3
cells at 20x in a particular experiment;

Fig. 16 illustrates 4 panels for each marker for a plurality of OVCAR-3 cells at 40x in a particular experiment;

Fig. 17 illustrates a representative input for a morphometric analysis program in a particular embodiment according to the present invention; and

5 Figs. 18-19 illustrate examples of the generation of pseudo-sequences and clustering in a particular embodiment according to the present invention.

Fig. 20 is a block diagram for a first research scenario;

Fig. 21 is a block diagram for a second research scenario; and

Fig. 22 is a block diagram for a third research scenario.

10 Reference numbers refer to the same or equivalent parts of the invention throughout the several figures of the Drawing.

DETAILED DESCRIPTION OF THE INVENTION

According to the present invention, techniques for determining information about manipulated cells or substances based upon living, fixed, or dead cell structures or portions of cells are provided. In an exemplary embodiment, the present invention provides a novel system including computer codes coupled to a database and user interfaces for acquiring, storing and retrieving such information. Other embodiments provide a novel image capturing system for providing digitized representations of live and dead cell structures or the like.

Fig. 1 is a simplified system diagram 10 of a cellular knowledge-based system according to an embodiment of the present invention. This diagram is merely an example and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize other variations, modifications, and alternatives. The present system 10 includes a variety of elements such as a computing device 13, which is coupled to an image processor 15 and is coupled to a database 21. The image processor receives information from an image capturing device 17, which image processor and image capturing device are collectively referred to as the imaging system herein. The image capturing device obtains information from a plate 19, which includes a plurality of sites for cells. These cells can be biological cells that are living, fixed, dead, cell fractions, cells in a tissue, and the like. The computing device retrieves the information, which has been digitized, from the image processing device and stores such information into the database. A user interface device 11, which can be a personal computer, a work station, a network computer, a personal digital assistant, or the like, is coupled to the computing device.

Fig. 1A is a simplified diagram of a database system 1000 according to an embodiment of the present invention. This diagram is merely an example and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize many other variations, modifications, and alternatives. Database system 1000 includes a variety of techniques for processing images from biological cells, e.g., fixed, living, and dead cells, and cell portions. As shown, images are acquired 1001. These images can be from a single frame or multiple frames. As merely an example, an image processing system may analyze such images. One example of

such an image processing system is described below, but should not be construed as limiting certain claims.

In a specific embodiment, cell samples are manipulated using a compound (e.g., substance, drug). The cell samples are imaged for a simple portion or portions, e.g., manipulated cell substructure, manipulated spatial feature of cell, cell density. Image processing techniques are used to extract the feature or features from the image or images. The features can be an independent or a dependent set of cell characteristics (which may be predominately visual) including, for example, count, area, perimeter, length, breadth, fiber length, fiber breadth, shape factor, elliptical form factor, inner radius, outer radius, mean radius, equivalent radius, equivalent sphere volume, equivalent prolate volume, equivalent oblate volume, equivalent sphere surface, average intensity, total intensity, optical density, radial dispersion, texture difference, and others. Each of these features corresponds to a similar manipulation by a compound. Each manipulation forms a new set of features, which are identifiable to the compound. Once each set of features has been extracted, the feature set is populated into a database. Accordingly, the database includes many sets of features, where each set corresponds to a different manipulation for a selected cell. Each set of features corresponding to a manipulation provides a descriptor, which is also stored in the database. The descriptor is a "finger print" including each feature for the manipulation. Each descriptor may be unique, or may have similarities to other descriptors or may even be the same as other descriptors for known and unknown manipulations.

The present system retrieves features, which we define as simple features herein, and forms composite features from them. More than one feature can be combined in a variety of different ways to form these composite features. In particular, the composite feature can be any function or combination of a simple feature and other composite features. The function can be algebraic, logical, sinusoidal, logarithmic, linear, hyperbolic, statistical, and the like. Alternatively, more than one simple feature can be combined in a functional manner (e.g., arithmetic, algebraic). As merely an example, the composite feature equals a sum of feature 1 and feature 2, where these features correspond to the same manipulation. Alternatively, the composite feature equals feature 1 divided by feature 2. Alternatively, the composite feature equals feature 1 minus feature 2. Alternatively,

the composite feature equals a constant times feature 1 plus feature 2. Of course, there are many ways that the composite feature can be defined. The present system also stores 1017 these features in the database. The composite features can also be further combined with simple features. Once these features are defined as descriptors, they are stored 1019 in the database.

Fig. 1B is a simplified diagram of a database system engine 2000 according to an embodiment of the present invention. This diagram is merely an example and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize many other variations, modifications, and alternatives. The engine can be implemented into the present database for populating, searching, and predicting compound or cell characteristics. As merely an example, engine 2001 includes an input/output module 2008. The input/output module is used to input and output information from the database. The information includes, among others, a plurality of feature sets, which correspond to many manipulations. Additionally, the information includes descriptors, which each corresponds to a set of features from the manipulation. The database also has a population module, which is used to configure the features based upon an entity relationship, which has been predetermined.

The database engine also has other modules. In particular, the database has a transcription module, which transfers a preselected set of features and creates a descriptor from them. The transcription module can be used to take a known compound, which has features, to transcribe them into a descriptor. Alternatively, the transcription module can be used to take an unknown compound, which has features, to transcribe them into a descriptor. These descriptors are provided into the database for subsequent use. Finally, the database engine has a prediction module, which can be used to potentially predict a property (e.g., mechanism of action) of an unknown compound. Here, the unknown compound is provided with a descriptor, but the property of the compound is unknown. In one embodiment, the prediction module compares a descriptor of an unknown compound with the many descriptors of known compounds, which were in the populated database. Depending upon the matching criteria, the prediction module will attempt to uncover one or more descriptors of known compounds. Once the prediction module finds the descriptors of the known compounds based upon the descriptor for the unknown compound, it identifies a potential property of such unknown compound for analysis and review. Here, it is

believed that certain features of the known compound, which are similar to those features of the unknown compound may uncover a property to the unknown compound. Details of the present software engine are described more fully below.

Fig. 2 is a simplified block diagram 20 of a cellular knowledge-based system according to an alternative embodiment of the present invention. This diagram is merely an example and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize other variations, modifications, and alternatives. Like reference numerals are used in the present diagram as the previous diagram for easy cross-referencing, but are not intended to be limiting in any manner.

10 The present diagram 20 includes a variety of elements such as a processor 13 or computing device coupled to a database 11. The processor can be used for retrieving and storing information from the database. The system also includes a plurality of system elements, such as a cleaner 23, a dispenser 25, and an image capturing system 27, which are also coupled to the database in some embodiments. These elements can

15 be coupled to each other through a network or the like. As merely an example, the network can be a NetWare™ network from Novell Corporation or an internet network or the Internet but can also be others and any combination thereof. The system also has an output device 31, which can be used to output information from the database, processor, or other system elements. Details of these elements are described more

20 fully below in reference to the Figs.

Figs. 3-5 are simplified drawings of system elements according to embodiments of the present invention. These diagrams are merely examples and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize other variations, modifications, and alternatives. As merely an example,

25 Fig. 3 is a simplified diagram of a processor or computing device 13. The computing device 13 includes a bus 112 which interconnects major subsystems such as a central processor 114, a system memory 116 (e.g., random access memory), an input/output ("I/O") controller 118, an external device such as a display screen 124 via a display adapter 126, a keyboard 132 and a mouse 146 via an I/O controller 118, a SCSI host

30 adapter (not shown), and a floppy disk drive 136 operative to receive a floppy disk 138.

The computing device has other features. Storage Interface 134 may act as a storage interface to a fixed disk drive 144 or a CD-ROM player 140 operative

to receive a CD-ROM 142. Fixed disk 144 may be a part of computing device or may be separate and accessed through other interface systems. A network interface 148 may provide a direct connection to a remote server via a telephone link or to the Internet. Network interface 148 may also connect to a local area network ("LAN") or other network interconnecting many computer systems. Many other devices or subsystems (not shown) may be connected in a similar manner. Also, it is not necessary for all of the devices shown in Fig. 3 to be present to practice the present invention, as discussed below. The devices and subsystems may be interconnected in different ways from that shown in Fig. 3. The operation of a computer system such as that shown in Fig. 3 is readily known in the art and is not discussed in detail in this application. Computer code to implement the present invention, may be operably disposed or stored in computer-readable storage media such as system memory 116, fixed disk 144, CD-ROM 140, or floppy disk 138. The computer code can be organized in terms of processes or modules, depending upon the application. That is, the computer code can include a prediction module, a translation module, or other modules to carryout the functionality described herein, as well as others.

Figs. 4 and 5 are simplified diagrams of an imaging system 200 according to an embodiment of the present invention. As shown, the imaging system 200 includes a variety of features such as housing 203, which holds a stage assembly 204. The stage assembly includes an x-stage movement element 206, which is along an x-direction, and a y-stage movement element 207, which is along a y-direction. The imaging system also includes a z-direction movement element, which is perpendicular to the x-y plane. The z-direction movement motor can be attached to the stage, or to the objective nosepiece by way of the microscope housing, or as an external motor between the objective and the microscope housing. The stage can align in any one of the directions to an accuracy of one micron and less, or one-half micron and less, or one-quarter micron and less, depending upon the embodiment.

The stage holds a plate 202 or cell holder, which houses one of a plurality of samples. The plate includes a spatial array 209 of process sites. Each of the process sites can include a plurality of cells and solutions depending upon the embodiment. Each of the sites can carry a sufficient amount of solution to prevent substantial evaporation of the sample during processing in some embodiments. In embodiments for large scale analysis, the plate includes at least 96 sites, or more than

or equal to 384 sites, or more than or equal to 1,536 sites. The plate bottom is transparent and thin, which allows light to pass through the sample. Additionally, the plate is made of a suitable chemical resistant material. As merely an example, the plate can be either a 96, or 384, or 1536 or other formats from places such as Becton Dickinson of Franklin Lakes, NJ, or Corning Science Products of Corning, NY. In a preferred embodiment, the plate is a Corning Costar black-walled 96 well plate catalog #3904 from Corning Science Products of Corning, NY, but should not be limited to these in some applications, but can be others.

Also shown is the condenser for the microscope 201, which can be used to collect phase, DIC, or bright field images of the cells. Images resulting from the illumination of the samples to fluorescence, phase, DIC, or bright field techniques are collected using an image capturing device 208, which captures an image or images of cells from the plate. In a specific embodiment, the microscope is an inverted configuration with the objectives on the bottom of the plate and the condenser disposed overlying an upper surface of the sites, while the image capturing device underlies the sites. Images captured by the imaging device, whether analogue or digital, are viewed by a monitor or other devices. The image capturing device can be any camera assembly such as a charge coupled device camera, which is known as a CCD camera, or other high resolution camera capable of capturing images from the sites. In a specific embodiment, the camera is an interline CCD camera which does not require an external shutter.

In a specific embodiment, the present imaging system can be any suitable unit that is flexible for automated image collection using multi-well plastic plates. The imaging system also should be adapted to collect high-resolution images of cells on plastic or glass plates, cell growth chambers, or coverslips. The system also can be used for imaging multiple cell markers in multiple imaging conditions. To accomplish this, the microscope system has a variety of elements such as a light source, a motorized excitation filter wheel and shutter, x-y-z-motorized stage, excitation and emission filters, Fluor phase and DIC objectives, motorized objective nosepiece, dichroic filters, motorized dichroic filter cubes, phase and DIC rings and prisms, CCD camera, and software control. As merely an example, the present imaging system can have components such as those listed in the Table below.

DESCRIPTION	MAKER	MODEL
Microscope	Zeiss	100M
(x-y) motorized stage	Prior	
Xenon lamp	Sutter	Lambda
Filter wheel	Sutter	Lambda-10
Microtitre Plate holder	Prior	500-H223R
Isolation Table	Kinetic Systems	9101-24-85
Objective Spacers	Polytec PI	P-721.90
Camera	Hamamatsu	C47-95
Computer	IBM	IntelliStation
Software	Metamorph	v.4
Objectives	Zeiss	Achroplan 10x/0.25 LD-Achroplan 20x/0.4 LD-Achroplan 40x/0.6

Table: Image Acquisition System Elements

5 In a specific embodiment, the present system has the following capabilities, which are not intended to be limiting.

Image acquisition

1) Ability to automatically acquire multi-wavelength images from multiple sites on one multi-well plate, to sequentially name image files, and to log any
10 imaging parameter information with image files.

2) Ability to link images with a larger database/spreadsheet of information.

3) Ability to automatically collect multiple plates by interfacing the imaging system with a robotic arm.

15

X-Y control

1) Ability to place 96, 384, or 1536 well plates onto microscope stage and move to each well sequentially.

2) Ability to return to each well and collect another round of images (multi-site time-lapse) or ability to collect rapid time-lapse information at each well (time-lapse of many wells).

3) Ability to collect a low magnification image, automatically
5 determine features which may be of interest, automatically change the objective to a higher magnification, and collect high magnification images of a fixed number of those identified cells in the sample.

4) Ability to collect multiple frames in each site.

10 Z control

1. Ability to auto-focus with substantially minimal damage to biological specimen or fluorophore.

2. Ability to auto-focus rapidly.

15 The present embodiment of the imaging system is shown by way of Figs. 5A and 5B. These diagrams are merely examples and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize other variations, modifications, and alternatives. The present imaging system 40 includes a variety of elements such as a microscope 41, which is preferably an epi-fluorescent microscope,
20 but can be confocal, multiphoton, or hybrid types. The microscope includes elements 41A, the motorized Z-axis; 41B, the motorized dichroic filter cube holder; and 41C, the motorized objective nosepiece. In one embodiment, the microscope is a Model 100M made by Zeiss. The microscope communicates to computer 51 through control lines 73, 75, and 76. The imaging system also has camera 50 coupled to controller
25 50A and computing device 51, which oversees and controls operations of the elements of the imaging system.

The present microscope includes drivers for spatially moving a stage in two dimensions, including an x-direction, a y-direction, and moving the objective nosepiece in a z-direction in a Cartesian coordinate system. The z-direction
30 movement is provided using a fast z-motor, which can make z-direction adjustments within a predetermined time. The z-direction movement generally provides for focussing of the sample to the camera. The focussing occurs within the predetermined time of preferably ten seconds and less, or five seconds and less, or one

second and less, depending upon the embodiment. As merely an example, the z-motor or positioner can be a model PIFOC objective nanopositioner made by a company called Physik Instrumente of Waldbronn, Germany, but also can be others. The z-motor couples to computer 51 through line 63, which may also include a
5 controller. Depending upon the embodiment, a second z-motor 41A connected to the computer 51 by line 73 may be used to keep the z-motor 42 in the center of its travel. Alternatively, in other embodiments the stage could be provided with a z-motor allowing for movement of the stage in the z-direction.

The present stage also includes an x-y stage 43. The x-y stage moves
10 plate 59, e.g., 96 site, 384 site, 1536 site. The x-y stage moves plate in an x-y spatial manner. The stage has an accuracy or repeatability of about 1 micron and less, or about 2 microns and less. The stage can move in a continuous manner or a stepped manner. The stage also can move up to 30 mm/sec. or faster. The stage also can move 1 mm/sec. and less, depending upon the embodiment. The stage can also step
15 0.1 micron and less or 1 micron and less, as well as other spatial dimensions. The stage can be one such as a Proscan Series made by Prior Scientific of Rockland, MA but can also be others. The stage is controlled via control line 61 through controller 43A, which couples to computer 51 through control line 65.

The stage includes plate holder 44. The plate holder can hold a single
20 plate. In other embodiments, plate holder can also hold multiple plates. The plate holder can use mechanical, electrical, fluid, vacuum and other means for holding the plate or plates. The plate holder also is sufficiently stable for securing the plate. As merely an example, the plate holder is a Model 500-H223R made by Prior Scientific of Rockland, MA. In some embodiments, the plate holder may need adjustment in
25 the z-direction to provide for a desirable focus of a sample on a plate. In these embodiments, the plate holder is supported by spacers 45 or a plurality of stage pins, which mechanically elevate the plate holder in the z-direction. These pins are generally made of a suitable material for supporting such plate holder and also are sufficiently resistant to chemicals and the like.

30 In some embodiments, the entire imaging system is placed on an isolation table 49. The isolation table is disposed between the microscope and support structure. The isolation table is designed to prevent excessive vibration of the plate. The isolation table is made of a suitable material such as steel and honeycomb but can

be others. The table has a thickness of about 8 inches or preferably less than about 24 inches. In one embodiment, the table is Model 9101-24-85 made by Kinetic Systems of Boston, MA.

The imaging system also has a lamp or illumination assembly 62. The lamp assembly provides for a light source (See reference letter B) to a plurality of elements in the imaging system. For easy reading, the light path is defined by the dotted lines, which are not intended to be limiting. The lamp assembly has a variety of elements such as a Xenon lamp 46. The Xenon lamp provides light at about 320 to 700 nanometers (Prefocused). The Xenon lamp is 175 or 300 Watts. As merely an example, the lamp can be a Lambda Model made by Sutter Instrument Company of Novato, CA.

Referring to Fig. 5B, the lamp assembly also has a cold mirror 58, an excitation filter wheel 48, excitation filter(s) 55, and an excitation light shutter 57. As shown, light is derived from the Xenon lamp, reflects off of the cold mirror 58, traverses through the excitation filter or filters 55, and is controlled by the excitation light shutter 57. The lamp assembly has filter wheel 48, which houses one of a plurality of filters, including excitation filters. The shutter and filter wheel are controlled via control lines 67, which are coupled to a computer 51 or other type of computing device. The control lines 67 are coupled through controller 57A (for element 57) and controller 48A (for element 48) via control line 69 to computer 51.

Preferably, light traverses from the lamp assembly through a light guide 47 to illuminate features within the plate. The light guide is suitably selected to have a flexible member, which can be used to place lamp source at a remote location away from the imaging device. The flexible member substantially keeps any vibration from the lamp assembly away from the imaging device. In some embodiments, the member is at least 1 foot away from the imaging device. The light guide is a guide, which is a flexible hose-type sleeve. The sleeve is filled with a liquid such as an aqueous solution containing chloride or phosphate. A thin layer may be formed on the inside of the sleeve. The layer can be a containing tetrafluoroethylene and hexafluoropropylene, or containing tetrafluoroethylene and perfluoromethyl vinyl ether, or tetrafluoroethylene and perfluoropropyl vinyl ether. An example of such a light guide is described in International Application No. WO/98/38537 filed February 29, 1997, and assigned to NATH, Gunther. The liquid

light guide has less than about 30% transmission loss of the light at a remote location such as the imaging system.

Light is derived from the lamp assembly and directs off of filter 56, which directs the light upward. Filter 56 can be a dichroic and emission filter, as well as others. The light traverses through microscope nosepiece 41C, and traverses through objective spacers 54. An objective 53 magnifies the light toward a predetermined point on the plate 59. The objective can be, for example, made by Zeiss of Jena, Germany, as well as other companies. The objective can be one of a plurality including 1X, 10X, 20X, 40X, and others, depending upon the application. Magnification can be further expanded or contracted by intermediate optics between the objective and the camera. Selection of filter or filters is controlled by computer 51 via control line 75.

The camera 50 captures an image of cells from plate 59. The image is obtained from light scattering off of cells or portions of cells in the plate through objective 53, through objective spacers, through filters 56, which are captured at camera 50. In this preferred embodiment, the camera is a digital camera, but can be an analogue camera. The digital camera is a CCD camera, which has 1280 by 1024 pixels, or more or less. The pixels can be 6.7 microns in dimension or more or less. The camera preferably is substantially free from an external shutter to quickly capture a plurality of images of cells from the plate. The camera is controlled via control line 71 through controller 50A, which connects to computer 51 through control line 70. The present invention can also include other types of image acquisition devices selected from at least an epifluorescence, a confocal, a total-internal reflection, a phase, a Hoffman, a bright field, a dark field, a differential interference contrast, an interference reflection, or multi-photon illumination device.

The present imaging system stores images on a high density memory device 60. The high density memory device is preferably optical, but can also be magnetic. The high density memory device can be any suitable unit that is capable of storing a plurality of images from a plurality of sites in the plate. The memory device can be a compact disk, which would generally use a compact disk burner or the like. Depending upon the embodiment, the high density memory device is used to archive the images that are captured from the camera in the imaging system. Further details

of the imaging system can be found throughout the present specification, and more particularly below.

As merely an example, the present invention can be implemented using the following sequence of steps, which have been described in a journal entry form.

- 5 Here, images are opened and objects are identified based on a background value that has been edited in starting image acquisition. Information is maintained in a spreadsheet or other database format, which has the following information for each object:

Image Name	Image Plane	Image Date and Time
Elapsed Time	Object #	Total area
Pixel area	Area	Hole area
Relative hole area	Standard area count	Perimeter
Length	Breadth	Fiber length
Fiber breadth	Shape factor	Ell. form factor
Inner radius	Outer radius	Mean radius
Average gray value	Total gray value	Optical density
Radial dispersion	Texture Difference Moment	EFA Harmonic 2, Semi-Major Axis
EFA Harmonic 2, Semi-Minor Axis	EFA Harmonic 2, Semi-Major Axis Angle	EFA Harmonic 2, Ellipse Area
EFA Harmonic 2, Axial Ratio	EFA Harmonic 3, Semi-Minor Axis	

10

After computations are done, the log file is saved. In particular, the file is saved in an appropriate place with an appropriate name.

In a specific embodiment, the present invention provides the following detailed example of journal entries, which should not limit the scope of the invention.

Set Up Sequential File Names	Interactive: user sets up prefix name and image storage directory
Open Data Log	Opens a DDE (Excel) File
Annotate Log File	Interactive: experimental information that will go into the first line of the log file of stage positions
Stage (Go to Origin)	Origin is set as the center of well A1
Stage (Move to Absolute Position)	Offset to upper left hand corner of well (1410, 1621)
Stage (Log Position)	-
Stage (Scan Wells)	User picks wells to scan: runs 3x3 image collection.jnl.

3X3 IMAGE COLLECTION.jnl

Stage (Scan)	Takes 9 images of well, -1600 motor steps apart from left to right 3 columns and 3 rows, runs FOCUS, COLLECT IMAGE, SAVE SEQUENTIAL FILE NAME.JNL.
--------------	---

5

FOCUS, COLLECT IMAGE, SAVE SEQUENTIAL FILE NAME.jnl.

Stage (Log Position)	Logs stage position of each image
ADC – Focus	Opens up the manual focusing window with whatever focus time is current set
Show Message and Wait	Interactive: user hits enter to continue when done focusing

ADC-Acquire from Digital Camera	Takes Hoechst image
Save Using Sequential File Names	
Close	Closes image window

START IMAGE ANALYSIS.jnl

Low Pass	3x3 convolution of already opened image
Low Pass	3x3
Show Region Statistics	Interactive: Show entire image statistics. Calculate background subtraction value for step 4. by: INTENSITY Average + INTENSITY Std. Dev.
Arithmetic	Interactive: User inputs subtraction value from 3. into the constant Value field
Threshold image	Creates threshold 1 unit above 0 to 4096
Integrated Morphometry – Load State	Loads Start Image Analysis.ima Classifier 100 < area < 200000
Integrated Morphometry – Measure	Interactive: Shows area summary information about all objects. The average number is used as the Standard Area in 8.
Object Standards - Set Object Standards	Interactive: User inputs average area value from 7. into Standard Area box to be used by automated IMA for all images

IMA OBJECTS.jnl

Low Pass	3x3 convolution
Low Pass	3x3 convolution
Arithmetic	This background subtraction value needs to be manually entered into this journal from the value determined in START IMAGE ANALYSIS.jnl step 3
Threshold Image	1 unit above 0
Integrated Morphometry – Load State	Hoecsht.IMA Classifier 200 < area < 200000
Integrated Morphometry – Measure	Measures statistical info for all objects
Run Journal	Runs log obj and sum data.jnl

Log obj and sum data.jnl

Integrated Morphometry – Log Data	Logs object data into Sheet 1
Integrated Morphometry – Log Data	Log summary data into Sheet 2

5

COLLECT AUTOMATED IMA DATA IN ONE SPREADSHEET.jnl

Run Journal	Runs OPEN OBJECT LOG DDE FILE.JNL
Loop for all Images in a Directory	Loops IMA OBJECTS.jnl
Close Summary Log	
Close Object Log	User must manually save Excel spreadsheet

OPEN OBJECT LOG DDE FILE.jnl

Open Object Log	Opens a DDE object log into sheet 1 of an Excel spreadsheet
Open Summary Log	Opens a summary log into sheet 2

COLLECT AUTOMATED IMA DATA IN ONE SPREADSHEET 16 BIT IMAGES.jnl

Arithmetic	Interactive: Opens Arithmetic window for user to input background subtraction level from START IMAGE ANALYSIS.jnl step 3
Run Journal	Runs OPEN OBJECT LOG DDE FILE.JNL
Loop for all Images in a Directory	Interactive: Runs IMA OBJECTS 16 bit.jnl. User picks directory from which to choose.

5

IMA OBJECTS 16bit.jnl

Low Pass	3x3 convolution
Low Pass	3x3 convolution
Copy to 8-bit Image	No autoscale, to new untitled image
Save Using Sequential File Name	Saves 8bit image using previously defined Sequential File names.
Arithmetic	This background subtraction value needs to be manually entered into this journal from the value determined in START IMAGE ANALYSIS 16 TO 8 BIT.jnl step 5
Threshold Image	1 unit above 0 to 255

Integrated Morphometry – Load State	Hoechst. IMA Classifier 200 < area < 200000
Integrated Morphometry – Measure	Measures statistical info for all objects
Run Journal	Runs log obj and sum data.jnl

START IMAGE ANALYSIS 16 to 8 BIT.jnl

Copy to 8-bit Image	No autoscale, to new untitled image
Close	Closes 16 bit image
Low Pass	3x3 convolution
Low Pass	3x3 convolution
Show Region Statistics	Interactive: Show entire image statistics. Calculate background subtraction value for step 6. by: INTENSITY Average + INTENSITY Std. Dev.
Arithmetic	Interactive: User inputs subtraction value from 5. into the constant Value field
Threshold image	Creates threshold by 1 unit above 0 to 255
Integrated Morphometry – Load State	Loads Start Image Analysis.ima Classifier 100 < area < 200000
Integrated Morphometry – Measure	Interactive: Shows area summary information about all objects. The average number is used as the Standard Area in 10.
Object Standards - Set Object Standards	Interactive: User inputs average area value from 9. into Standard Area box to be used by automated IMA for all images

IMA OBJECTS WITH NEW LOG FILE.jnl

Run Journal	OPEN OBJECT LOG DDE FILE.JNL
Run Journal	IMA OBJECTS.jnl
Close Summary Log	
Close Object Log	User must manually save every Excel spreadsheet generated.

INTERACTIVE IMA OBJECTS.jnl

Threshold Image	User manually sets threshold
Integrated Morphometry – Load State	Hoechst.IMA Classifier 200 < area < 200000
Integrated Morphometry – Measure	Objects
Integrated Morphometry – Log Data	Into open object.log file

5

COLLECT INTERACTIVE IMA DATA.jnl

Close Object Log	
Open Object Log	Interactive
Annotate Log File	Interactive: experimental information that will go into the first line of the object log file
Loop for all Images in Directory	Runs INTERACTIVE IMA OBJECTS.jnl

CHANGE FILTER, COLLECT IMAGE, SAVE SEQUENTIAL FILE
NAME.jnl

Stage (Log Position)	
ADC-Focus	

Show Message and Wait	Interactive – user presses Enter when done focusing
ADC – Acquire from Digital Camera	Hoechst
Save Using Sequential File Name	
Close	Close open image

COLLECT HOECHST AND FITC.jnl

Run Journal	FOCUS, COLLECT IMAGE, SAVE SEQUENTIAL FILE NAME.JNL
Run Journal	CHANGE FILTER, COLLECT IMAGE, SAVE SEQUENTIAL FILE NAME.jnl

3X3 IMAGE COLLECTION HOECHST FITC.jnl

Stage (Scan)	COLLECT HOECHST AND FITC.jnl
--------------	------------------------------

5

AUTOMATED 3X3 IMAGE COLLECTION HOECHST FITC.jnl

Set Up Sequential File Names	Interactive: user sets up prefix name and image storage directory
Open Data Log	Excel DDL files
Annotate Log File	Interactive: experimental information that will go into the first line of the log file of stage positions
Stage (Go to Origin)	Origin is set as the center of well A1
Stage (Move to Absolute Position)	Offset to upper left hand corner of well (1410, 1621)

Stage (Log Position)	
Stage (Scan Wells)	Interactive: user picks wells to scan: runs 3X3 IMAGE COLLECTION HOECHST FITC.jnl

AUTOMATED IMAGE COLLECTION.jnl

Set Up Sequential File Names	Interactive: user sets up prefix name and image storage directory
Open Data Log	Opens a DDE (Excel) File
Annotate Log File	Interactive: experimental information that will go into the first line of the log file of stage positions
Stage (Go to Origin)	Origin is set as the center of well A1
Stage (Log Position)	
Stage (Scan Wells)	Interactive: user picks wells to scan: runs FOCUS, COLLECT IMAGE, SAVE SEQUENTIAL FILE NAME.JNL. Well to well travel = (-9035, -9035)

5

STARTUP.jnl

Install and Configure Devices	Open Stage Meta Devices
Set Live Video Channel	

Preferences	<u>Measure Objects</u> : Draw failed classifier objects, Exclude objects that touch the edge of the image, Enable Elliptical Fourier Parameters, turn off Warn users when measurement data will be erased <u>Image Saving</u> : Save Tiff/stk using LZW compression <u>Image Windows</u> : Use transparent thresholds.
Configure Default Paths	C:\Metamorph Data C:\Metamorph Data\Common Settings
Load Journal Taskbar	Common.JTB

Nested Journals

Automated 3x3 Image Collection

- 5 *Loop* 3x3 image collection
 Loop focus, collect image, save sequential file name

Automated 3x3 image collection Hoechst FITC

- 10 *Loop* 3x3 image collection Hoechst FITC
 loop Collect Hoechst and FITC
 focus, collect image, save sequential file name
 change filter, collect image, save sequential file name

Automated image collection

- 15 *Loop* focus, collect image, save sequential file name

Collect automated IMA data in one Spreadsheet

Open object log DDE file

Loop IMA objects

Log obj and sum data

Collect automated IMA data in one spreadsheet 16 bit images

5 Open object log DDE file

Loop IMA objects 16 bit

Log obj and sum data

Although the above has been generally described in terms of a specific
10 user interface and software code, other user interfaces and code can also be used. One
of ordinary skill in the art would recognize many other variations, alternatives, and
modifications.

Fig. 6 is a simplified diagram 600 of a cleaning and dispensing system
according to an embodiment of the present invention. This system 600 includes a
15 variety of elements such as a dispensing head 609, which is coupled to a plurality of
pipettes 601. The pipettes input and output fluids or solutions from plate 603. The
plate has a plurality of sites, each of which can be used to input cells or a combination
of cells and solution. The system also has elements to house solutions 605, which are
used to manipulate cell samples in the plate. The dispensing head is supported
20 through a support member 607, which is sufficiently rigid to allow for movement of
the head. The dispenser is coupled to the present system in a mechanical and
electrical manner, which provides for a fully integrated system for providing cell
samples to the imaging system according to the present invention.

Fig. 7A illustrates a representative block flow diagram of simplified
25 process steps of a method for determining properties of a manipulation based upon
effects of the manipulation on one or more portions of one or more cells in a
particular embodiment according to the present invention. This diagram is merely an
illustration and should not limit the scope of the claims herein. One of ordinary skill
in the art would recognize other variations, modifications, and alternatives. In step
30 700, one or more samples of cells can be provided. These cells can be live, dead, or
fixed cells, or cell fractions. The cells also can be in one of many cell cycle stages,
including G0, G1, S, G2 or M phase, M phase including the following cell cycle
stages: interphase, prophase, prometaphase, metaphase, anaphase, and telophase.

Cell components tracked in presently preferable embodiments can include proteins, protein modifications, genetically manipulated proteins, exogenous proteins, enzymatic activities, nucleic acids, lipids, carbohydrates, organic and inorganic ion concentrations, sub-cellular structures, organelles, plasma membrane, adhesion complex, ion channels, ion pumps, integral membrane proteins, cell surface receptors, G-protein coupled receptors, tyrosine kinase receptors, nuclear membrane receptors, ECM binding complexes, endocytotic machinery, exocytotic machinery, lysosomes, peroxisomes, vacuoles, mitochondria, Golgi apparatus, cytoskeletal filament network, endoplasmic reticulum, nuclear membrane, proteosome apparatus, chromatin, nucleolus, cytoplasm, cytoplasmic signaling apparatus, microbe specializations and plant specializations.

The following table illustrates some markers and cell components commonly used by embodiments according to the present invention. Other markers can be used in various embodiments without departing from the scope of the invention.

Cell component	Marker	Disease State
Plasma membrane (including overall cell shape)	Carbocyanine dyes Phosphatidylserine Various lipids Glycoproteins	Apoptosis-Cancer Apoptosis-Neural degenerative Ds
Adhesion complexes	Cadherins Integrins Occludin Gap junction ERM proteins CAMs Catenins Desmosomes	Thrombosis Metastasis Wound healing Inflammatory Ds Dermatologic Ds
Ion Channels and Pumps	Na/K Atpase Calcium channels Serotonin reuptake pump CFTR	Cystic fibrosis Depression Congestive Heart Failure Epilepsy

G coupled receptors	β adrenergic receptor Angiotensin receptor	Hypertension Heart Failure Angina
Tyrosine kinase receptors	PDGF receptor FGF receptor IGF receptor	Cancer Wound healing Angiogenesis Cerebrovascular Ds
ECM binding complexes	Dystroglycan Syndecan	Muscular Dystrophy
Endocytotic machinery	Clathrin Adaptor proteins COPs Presenilins Dynamin	Alzheimer's Ds
Exocytotic machinery	SNAREs Vesicles	Epilepsy Tetanus Systemic Inflammation Allergic Reactions
Lysosomes	Acid phosphatase Transferrin	Viral diseases
Peroxisomes/Vacuoles		Neural degenerative Ds
Mitochondria	Caspases Apoptosis inducing factor F1 ATPase Fluorescein Cyclo-oxygenase	Apoptosis Neural degenerative Ds Mitochondrial Cytopathies Inflammatory Ds
Golgi Apparatus	Lens Culinaris DiOC6 carbocyanine dye COPs	

Cytoskeletal Filament Networks	Microtubules Actin Intermediate Filaments Kinesin, dynein, myosin Microtubule associated proteins Actin binding proteins Rac/Rho Keratins	Cancer Neural degenerative Ds Inflammatory Ds Cardiovascular Ds Skin Ds
Endoplasmic Reticulum	SNARE PDI Ribosomes	Neural degenerative Ds
Nuclear Membrane	Lamins Nuclear Pore Complex	Cancer
Proteosome Apparatus	Ubiquityl transferases	Cancer
Chromatin	DNA Histone proteins Histone deacetylases Telomerases	Cancer Aging
Nucleolus	Phase markers	
Cytoplasm	Intermediary Metabolic Enzymes BRCA1	Cancer
Cytoplasmic Signaling Apparatus	Calcium Camp PKC pH	Cardiovascular Ds Migraine Apoptosis Cancer
Microbe Specializations	Flagella Cilia Cell Wall components: Chitin synthase	Infectious Ds

Plant specializations	Choloroplast Cell Wall components	Crop Protection
-----------------------	--------------------------------------	-----------------

Then, in a step 702, one or more samples of the manipulation can be provided to the cells. Manipulations can comprise one or any combination of chemical, biological, mechanical, thermal, electromagnetic, gravitational, nuclear, or temporal factors, for example. For example, manipulations could include exposure to chemical compounds, including compounds of known biological activity such as therapeutics or drugs, or also compounds of unknown biological activity. Or exposure to biologics that may or may not be used as drugs such as hormones, growth factors, antibodies, or extracellular matrix components. Or exposure to biologics such as infective materials such as viruses that may be naturally occurring viruses or viruses engineered to express exogenous genes at various levels. Bioengineered viruses are one example of manipulations via gene transfer. Other means of gene transfer are well known in the art and include but are not limited to electroporation, calcium phosphate precipitation, and lipid-based transfection. Manipulations could also include delivery of antisense polynucleotides by similar means as gene transfection. Other genetic manipulations include gene knock-outs or gene mutations. Manipulations also could include cell fusion. Physical manipulations could include exposing cells to shear stress under different rates of fluid flow, exposure of cells to different temperatures, exposure of cells to vacuum or positive pressure, or exposure of cells to sonication. Manipulations could also include applying centrifugal force. Manipulations could also include changes in gravitational force, including sub-gravitation (the preferred embodiment in outer space). Manipulations could include application of a constant or pulsed electrical current. Manipulations could also include irradiation. Manipulations could also include photobleaching which in some embodiments may include prior addition of a substance that would specifically mark areas to be photobleached by subsequent light exposure. In addition, these types of manipulations may be varied as to time of exposure, or cells could be subjected to multiple manipulations in various combinations and orders of addition. Of course, the type of manipulation used depends upon the application.

Then, in a step 704, one or more descriptors of a state in the portions of the cells in the presence of the manipulation can be determined using the images

collected on the imaging system. Descriptors can comprise scalar or vector values, representing quantities such as area, perimeter, dimensions, intensity, gray level, aspect ratios, and the like. Other types of descriptors include, but are not limited to, one or any combination of characteristics such as a cell count, an area, a perimeter, a length, a breadth, a fiber length, a fiber breadth, a shape factor, a elliptical form factor, an inner radius, an outer radius, a mean radius, an equivalent radius, an equivalent sphere volume, an equivalent prolate volume, an equivalent oblate volume, an equivalent sphere surface area, an average intensity, a total intensity, and an optical density. These descriptors can be average or standard deviation values, or frequency statistics from the descriptors collected across a population of cells. These descriptors can be further reduced using other methods such as principal component analysis and the like. In some embodiments, the descriptors include features from different cell portions or cell types. That is, a first feature can be from a nuclei and a second feature is from another cell structure such as Golgi apparatus, mitochondria, spacing between cell structures or cells themselves, as well as many others.

A presently preferable embodiment uses descriptors selected from the following table. Other descriptors can also be used without departing from the scope of the invention.

Name of Parameter	Explanation/Comments
Count	Number of objects
Area	
Perimeter	
Length	X axis
Width	Y axis
Shape Factor	Measure of roundness of an object
Height	Z axis
Radius	
Distribution of Brightness	
Radius of Dispersion	Measure of how dispersed the marker is from its centroid
Centroid location	x-y position of center of mass
Number of holes in closed objects	Derivatives of this measurement might include, for

	example, Euler number (= number of objects - number of holes)
Elliptical Fourier Analysis (EFA)	Multiple frequencies that describe the shape of a closed object
Wavelet Analysis	As in EFA, but using wavelet transform
Interobject Orientation	Polar Coordinate analysis of relative location
Distribution Interobject Distances	Including statistical characteristics
Spectral Output	Measures the wavelength spectrum of the reporter dye. Includes FRET
Optical density	Absorbance of light
Phase density	Phase shifting of light
Reflection interference	Measure of the distance of the cell membrane from the surface of the substrate
1,2 and 3 dimensional Fourier Analysis	Spatial frequency analysis of non closed objects
1,2 and 3 dimensional Wavelet Analysis	Spatial frequency analysis of non closed objects
Eccentricity	The eccentricity of the ellipse that has the same second moments as the region. A measure of object elongation.
Long axis/Short Axis Length	Another measure of object elongation.
Convex perimeter	Perimeter of the smallest convex polygon surrounding an object
Convex area	Area of the smallest convex polygon surrounding an object
Solidity	Ratio of polygon bounding box area to object area.
Extent	proportion of pixels in the bounding box that are also in the region
Granularity	
Pattern matching	Significance of similarity to reference pattern
Volume measurements	As above, but adding a z axis

Then, in a step 705, a database of cell information can be provided.

Next, in a step 706, a plurality of descriptors can be searched from a database of cell information in order to locate descriptors based upon one of the descriptors of the manipulation. Then, in a step 708, properties of the manipulation are predicted based
5 upon the properties of the located descriptors. Properties can comprise toxicity, specificity against a subset of tumors, mechanisms of chemical activity, mechanisms of biological activity, structure, adverse biological effects, biological pathways, clinical effects, cellular availability, pharmacological availability, pharmacodynamic properties, clinical uses and indications, pharmacological properties, such as
10 absorption, excretion, distribution, metabolism and the like.

In a particular embodiment, step 706 comprises determining matching descriptors in the database corresponding to a prior administration of the manipulation to the descriptors of the present administration of the manipulation. In a particular embodiment according to the present invention, combinations of
15 measurements of scalar values can provide predictive information. A database can be provided having one or more "cellular fingerprints" comprised of descriptors of cell-substance interactions of drugs having known mechanisms of action with cells. Such descriptors can be analyzed, classified, and compared using a plurality of techniques, such as statistical classification and clustering, heuristic classification techniques, a
20 technique of creating "phylogenetic trees" based on various distance measures between descriptors from various drugs. In this embodiment, numeric values for the descriptors can be used by comparison techniques. A phylogenetic tree can be created that illustrates a statistical significance of the similarity between descriptors for the drugs in the database. Because the drugs used to build the initial database are of
25 known mechanism, it can be determined whether a particular scalar value in a descriptor is statistically predictive. Finally, a compound descriptor with no known mechanism of action can be queried against the database and be statistically compared and classified among the drugs in the database that the compound most resembles.

In a particular embodiment, relationships between measured
30 morphological properties of images and physiological conditions can be determined. Relationships can include, for example, treatment of different cell lines with chemical compounds, or comparing cells from a patient with control cells, and the like. In a presently preferable embodiment, comparisons can be performed on acquired image

features. Some embodiments can comprise statistical and neural network - based approaches to perform comparisons of various features. The foregoing is provided as merely an example, and is not intended to limit the scope of the present invention. Other techniques can be included for different types of data.

- 5 In some embodiments, classification, clustering and other types of predictive data analysis can be performed on features extracted from cell images. In a presently preferable embodiment, statistical procedures for comparisons, classification and clustering are performed on data obtained from imaging cells.

Fragments of data preparation and pre-formatting (S language):

```
10 >tmp.frame <- Generic.Summary
    >names1 <- paste("Cell.line.5", tmp.names, sep=".")
    > by.compound.matrix <- as.matrix(arranged.by.compound)
```

- 15 Example of the code for principal component analysis (data preparation) using S language:

```
    all.data.princomp <- menuPrincomp(data =
    by.compound.matrix, scores = T, cor = "Correlation",
    na.action = T, print.short = T, print.importance = T,
    print.loadings = T, cutoff.loadings = 0.1,
    20 plot.screepplot = T, plot.loadings = T, plot.biplot = T,
    plot.biplot.choices = c(1,2), predict.p = F)
```

Example of clustering using a divisive hierarchical clustering algorithm:

```
25 > div.hier.2.manhattan.cluster$call
    diana(x = tmp.sum.by.comp, diss = F, metric =
    "manhattan",
    stand = T, save.x = T, save.diss = T)
```

- 30 Another embodiment utilizes existing tools for biological sequence similarity searches, classification, and phylogenetic analysis. In a particular embodiment, numbers in a numerical descriptor can be substituted by one or more of nucleic acid or amino acid codes according to a one of several sets of rules. Once

converted into a corresponding nucleotide or amino acid sequence representation, the fingerprints can be analyzed and compared using software and algorithms known in the art for genetic and peptide sequence comparisons, such as GCG, a product of Genetics Computer Group, with company headquarters in Madison WI. Select
5 embodiments comprising such approaches enable the use of a broad array of sophisticated algorithms to compare, analyze, and cluster gene and protein sequences. Many programs performing this task are known to those of ordinary skill in the art, such as for example, the PHYLIP (PHYlogeny Interference Package) a package of programs for inferring phylogenies (evolutionary trees) described in (Feldenstein, J.
10 1996 Methods Enzymol 266:418-427 and Feldenstein, J. 1981 J. Mol. Evol. 17(6):368-376).

Embodiments can perform such analysis based upon factors such as numerical value, statistical properties, relationships with other values, and the like. Further details of a step of manipulation are noted more particular below.

15 Fig. 7B illustrates a representative block flow diagram of simplified process steps for determining one or more descriptors of a state in the portions of the cells in the presence of the manipulation of step 704 of Fig. 7A in a particular embodiment according to the present invention. This diagram is merely an illustration and should not limit the scope of the claims herein. One of ordinary skill
20 in the art would recognize other variations, modifications, and alternatives. In a step 712, an image of a cell portion is obtained. In some embodiments, the cell portion is visualized with a fluorescently labeled marker that is specific for the portion or portions of interest. A cell portion can include, for example, one or more of the following: nuclei, Golgi apparatus, and other features. The cell portion may vary in
25 select embodiments according to the invention. Then, in a step 714, a digitized representation of the image obtained in step 712 is determined. In some embodiments, steps 714 and step 712 can comprise a single step. These embodiments use a digital imaging means such as a digital camera, to obtain a digital image of the target directly. Next, in a step 716, the digital representation of the image is
30 processed to obtain image features. Image features can include such quantities as area, perimeter, dimensions, intensity, aspect ratios, and the like. Then, in a step 718 descriptors can be determined from the image features. Descriptors can comprise scalar or vector quantities and can comprise the image features themselves, as well as

composed features, such as shape factor derived by a relationship $4\pi \cdot \text{area} / \text{perimeter}$, and the like. Descriptors can also comprise statistical quantities relating to feature characteristics across a population of cells, such as a standard deviation, and average, and the like.

5 In a preferred embodiment, cells can be placed onto a microscope, such as a Zeiss microscope, or its equivalent as known in the art. A starting point, named Site A01, is identified to the microscope. A plurality of exposure parameters can be optimized for automated image collection and analysis. The microscope can automatically move to a new well, automatically focus, collect one or more images, at
10 one or more wavelengths, move to a next well, and repeat this process for all designated wells in a multiple well plate and for multiple plates. A file having a size and an intensity distribution measurement for each color and rank for each well can then be created for the images acquired. Based on this information, a user or a computer can revisit sites of interest to collect more data, if desired, or to verify
15 automated analysis. In a presently preferred embodiment, image automatic focus and acquisition can be done using computer software controlling the internal Z-motor of the microscope. Images are taken using a 10x, 20x, or 40x air long working distance objectives. Sometimes multiple images are collected per well. Image exposure times can be optimized for each fluorescent marker and cell line. The same exposure time
20 can be used for each cell line and fluorescent marker to acquire data.

Fig. 7C illustrates a representative block flow diagram of simplified process steps for obtaining images of cell portions of step 712 of Fig. 7B in a particular embodiment according to the present invention. This diagram is merely an illustration and should not limit the scope of the claims herein. One of ordinary skill
25 in the art would recognize other variations, modifications, and alternatives. The method is generally outlined by the steps below:

- (1). In a step 720, a sample is provided to the imaging device. Samples can be provided in 96 well plates and the like. The sample may be loaded into a microscope, such as a Zeiss microscope or equivalent.
- 30 (2). In a step 722, a set of optical filters is selected to shine light of the appropriate wavelength to illuminate the first sample, which may be contained in a first well designated A01.

(3). In a step 724, an automatic focusing procedure is performed for the site. In a particular embodiment, the internal z-motor of the microscope which is attached to the objective nosepiece is used for automatic focusing of the microscope. In an alternative embodiments, the plate holding the samples is moved to perform
5 automatic focusing of the microscope, or focusing can be performed by moving optical components attached to the microscope and the like.

(4). In a step 726, images are collected for the site. Images can be collected for every color at every site. Present embodiments can provide images for up to four colors. However, embodiments are contemplated that can provide more
10 colors by using either a monochromator coupled with excitation filters which are on a filter wheel, or by digitally separating overlapping fluorophores. Those knowledgeable in the field will know that given calibration images of single fluorophores, a look-up table can be devised which will allow for the digital removal of fluorescence bleed-through of fluorescence which may occur in optical channels
15 other than the one for which that filter has been optimized in instances of using more than one fluorophore at once. Cell growth and density information is also collected. Cell density is determined by what percentage of the area being imaged is inhabited by cells. In some embodiments, imaging can be facilitated using one or more biosensors, molecules such as non-proteins, i.e., lipids and the like, that are
20 luminescently tagged. However, some embodiments can also use fluorescence polarization and the like. Fluorescence polarization is a homogeneous fluorescence technology where the excited state of the molecule lasts much longer than in normal fluorescence, taking seconds to minutes to reach equilibrium, obliterating the need to wash away fluorescence markers that are not specifically bound to a marker. Further,
25 embodiments can detect differences in spectral shifts of luminescent markers. Some fluorescence markers, such as Nile Red sold by Molecular Probes of Eugene, OR, will change its emission peak wavelength depending on its environment. One can detect these changes by monitoring the level of fluorescence at both wavelengths and reading out at ratio of the two.

30 (5). In a step 728, a determination is made whether more fields of view need to be taken for a particular color. If this is so, then processing continues at step 726 at a new site. Otherwise, processing continues with a decisional step 730.

Images can now be taken by repeating step 726. In a preferred embodiment 4 to 9 images are collected at each site.

(5). In a step 730, a determination is made whether more optical configurations need to be taken in order to obtain images for all differently-marked cell portions the sample. If this is so, then in a step 732 a new optical configuration is determined. Images for the new optical configuration can now be taken by repeating steps 726 and 728.

(6). In a decisional step 734, after all optical configurations and images for fields of view in a sample have been obtained, a determination is made whether any further samples remain to be analyzed. If so, a new sample is brought into view and processing continues with step 720. Otherwise, image processing is complete. In a presently preferable embodiment, image data can be stored on a CD ROM using a CD ROM burner, such as CRW4416 made by Yamaha of Japan. However, other mass storage media can also be used.

Fig. 7D illustrates a representative block flow diagram of simplified process steps for processing digitized representations of step 716 of Fig. 7B in a particular embodiment according to the present invention. This diagram is merely an illustration and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize other variations, modifications, and alternatives. The method is generally outlined by the steps below:

(1). In a step 740, a digitized image input is preprocessed. Preprocessing might include, but is not limited to, such operations as background subtraction, thresholding, smoothing, adoptive filtering, edge enhancements, contrast enhancements, histogram equalization. A particular combination of preprocessing steps can be applied to images in successive steps or in parallel to copies of the image.

A simplified example of a smoothing and background subtraction procedure in a MatLab language is presented in computer code below:

```
function Isubtracted = cmBackgrSubtr1(I,k)
% cmBackgrSubtr1(I,k) - simple flat background (=modal*k)
subtraction
% Y = cmBackgrSubtr1(I, k) - image Y is generated by
```



```

    % subtraction (with saturation) of modal pixel value of I
    multiplied by k
    % DEFAULT - k=1
    %
5   if (nargin == 1)
        k=1;
    end
    if (size(k)~=1)
        error('cmBackgrSubtrl: parameter k should be a number.
10  Exiting...');
    end

    %modpixnum = floor(size(I(:),1)/2);
    %sortedval = sort( double(I(:)) );
15  %modpixel = sortedval(modpixnum);
    modpixel = median(double(I(:)));
    bg = k*modpixel;

    Isubtracted = mmsubm( uint8(I), uint8(round(ones(
20  size(I))*k*modpixel )) );

```

An example of a procedure for thresholding in computer code (MatLab) is presented below:

```

function thresh = GetThreshByPerim1(I, M)
25 % GetThreshByPerim1(I) Finds optimal thresholding value
    for image I
    % N = GetThreshByPerim1(I) Finds thresholding value N for
    image I
    % N = GetThreshByPerim1(I, M) - tests threshold values up
30  to M
    % DEFAULT M = maximum pixel value in I
    % note that GetThreshByArea is significantly faster
    % finds a threshold value that causes the maximal change
    in the

```

```

% total perimeter of the objects (Russ ????)
% see Matlab_Auto_threshold1_1-23-99.doc for more details
% Note: works somewhat better on SMOOTH images (i.e.
medfilt2(I, [3 3]) two times

5
if (nargin == 0)
    error (strcat( mfilename, ' : at least one parameter
required')));
elseif (nargin == 1)
10    M = double(max(I(:)));    %test thresholds up to
    maximum pixel value in I
elseif (nargin > 2)
    error (strcat (mfilename, ' : too many parameters'));
end

15
if (size(M)>1)
    error (strcat(mfilename, ' : argument M should be a
number')));
end

20
Minval = double( min(I(:)));
step = 1;

%generate vertical vector perims with total perimeters of
25 objects at different
%threshold values
for i=Minval : step : M
    bwI = im2bw(I, i/255);
    prI = bwperim(bwI);
30    pr = sum(prI(:));
    if (exist('perims', 'var') == 0) %perims is yet
undefined
        perims = pr;
    else

```

```

        perims = cat(1, perims, pr);
    end
end

5  % vector prdiffs contains differences between successive
    perimeters
    prdiffs = diff(perims);
    mindecrease = min(prdiffs);
    minvalues = find(prdiffs == mindecrease);
10  index_of_mindecrease = minvalues(1);
    thresh = index_of_mindecrease + 1;

    % =====end GetThresh1=====

```

15 Thresholding provides a specific intensity, such that pixels darker than the threshold are deemed black, and pixels lighter than the threshold are considered white. The thresholded image can be processed using binary image processing techniques in order to extract regions.

(2). In a step 742+744, the digitized image input is subjected to object
20 identification. This can be accomplished by a variety of procedures, for example by thresholding or edge detection and subsequent morphological opening and closing. Edge detection can be accomplished by means of gradient-based or zero-crossing methods, such as Sobel, Canny, Laplassian, Perwitt, and other methods.

An example of object identification procedure based on Canny edge
25 detection (in MatLab language) is presented below:

```

function Imask = cmMaskDNA1(I);
% cmMaskDNA1 - generates binary mask for cell nuclei
through edge detection
30 % Imask = cmMaskDNA1(I)
% PARAMETERS
%   I - intensity image (grayscale)
% OUTPUT
%   Imask - BW image with objects from I

```

```

%
% For more details see Notebook Matlab_DNA_masking1_1-22-
99.doc
% Uses SDC Morphology Toolbox V0.7
5
if (nargin ~= 1)
    error('Wrong number of input parameters');
end
if (nargout ~= 1)
10    error('Wrong number of output parameters: one output
argument should be provided');
end

15  Imask = edge(I, 'canny');
    Imask = mmdil(Imask, mmsecross(1));
    Imask = mmero ( mmclohole(Imask,mmsecross(1)));
    Imask = mmedgeoff(Imask, mmsecross(1));
    % note that mmedgeoff this command removed FILLED OBJECTS
20  but not touching OUTLINES.
    % these outlines can be removed by filtering:
    Imask = medfilt2(Imask, [5 5]);

    %=====end cmMaskDNA1
25  =====

```

However, embodiments can also use other techniques, such as Fast Fourier Transforms (FFT) and the like as known in the art without departing from the scope of the present invention.

30 (3). In a step 746, a plurality of region features can be determined. For example, in a representative embodiment, image features can include such quantities as area, perimeter, dimensions, intensity, aspect ratios, and the like. Features not directly related to individual objects are also being extracted.

An example of a procedure for extraction of some of the features (MatLab language) is presented below:

```

function OData = cmGetObjectsData(I, Ilabel)
5  % cmGetObjectsData returns array measurements of objects
   in image "I" masked by "Ilabel"
   % EV 2-3-99; 2-10-99
   % OData = cmGetObjectsData(I, Ilabel) returns an array of
   morphological and intensity measurements
10  %   taken from a grayscale image "I". Objects are
   identified on a mask image Ilabel, usually
   %   created by bwlabel()
   % OUTPUT:
   % Each row in the output array OData represents
15  individual object
   % columns contain the following measurements:
   %
   %   1 - Index ("number" of an object);      8 -
   Solidity;
20  %   2 - X coordinate of the center of mass; 9 - Extent;
   %   3 - Y coordinate      "-"; 10 - Total
   Intensity;
   %   4 - Total Area (in pixels);      11 - Avg.
   Intensity;
25  %   5 - Ratio of MajorAxis/MinorAxis;      12 - Median
   Intensity;
   %   6 - Eccentricity;      13 - Intensity of
   20% bright pixel
   %   7 - EquivDiameter;      14 - Intensity of
30  80% bright pixel
   %
   % For details on morphological parameters see information
   on MatLab imfeature();

```

```
% Intensity parameters are either obvious or are
documented in comments in this file.

if (nargin ~= 2)
5   error ('function requires exactly 2 parameters');
end
if (nargout ~= 1)
    error ('function has 1 output argument (array X by
14)');
10 end

% finished checking arguments

% first collect morphological parameters in a structure
15 array:
ImStats = imfeature(Ilabel, 'Area', 'Centroid',
    'MajorAxisLength',...
    'MinorAxisLength', 'Eccentricity', 'EquivDiameter',
    ...
20 'Solidity', 'Extent', 8 );

% now convert it into array (matrix) while collecting
intensity data for each object:

25 %preallocate output array:
numobjects = size(ImStats, 1);
OData = zeros(numobjects, 14);
%now convert ImStats into array and add intensity data to
it
30 for k=1:numobjects
    OData(k, 1) = k;
    OData(k, 2) = ImStats(k).Centroid(1);
    OData(k, 3) = ImStats(k).Centroid(2);
    OData(k, 4) = ImStats(k).Area;
```

```

        OData(k, 5) = (ImStats(k).MajorAxisLength) /
        (ImStats(k).MinorAxisLength);
        OData(k, 6) = ImStats(k).Eccentricity ;
        OData(k, 7) = ImStats(k).EquivDiameter;
5       OData(k, 8) = ImStats(k).Solidity;
        OData(k, 9) = ImStats(k).Extent;

        % now collect and assign intensity parameters from
        image I
10
        object_pixels = find( Ilabel == k);
        object_area = size(object_pixels, 1); %same as total
        number of pixels in the object
        object_intensities = double(I(object_pixels)); %
15 need to convert to double to do math
        sorted_intensities = sort(object_intensities); %
        will need to get median, 20% and 80% pixels
        total_intensity = sum(object_intensities, 1);
        avg_intensity = total_intensity / object_area;
20 median_intensity = sorted_intensities( floor(
        object_area/2 ) + 1 );
        pix20 = sorted_intensities( floor(object_area*0.2)+1
        ) ; %brightest pixel among dimmest 20%
        pix80 = sorted_intensities( floor(object_area*0.8)+1
25 ) ;

        OData(k, 10) = total_intensity;
        OData(k, 11) = avg_intensity;
        OData(k, 12) = median_intensity;
30 OData(k, 13) = pix20; %brightest pixel among dimmest
        20%
        OData(k, 14) = pix80; %dimmest pixel among brightest
        20%
        end %for

```

```
%===== end function
cmGetObjectsData() =====
```

- 5 (4). In a step 748, quantitative descriptors, characterizing cell state are calculated based on the feature measurements extracted at step 746. For example, histogram distribution of intensities of cell nuclei provides information about the population cell cycle stages.

In a particular embodiment according to the present invention, data analysis techniques for describing the fluorescence patterns of cell portions in multiple cell lines in the presence and absence of compounds are provided. Automated image analysis techniques can include determining one or more regions from around nuclei, individual cells, organelles, and the like, called "objects" using a thresholding function. Objects that reside on the edge of an image can be included or excluded in various embodiments. An average population information about an object can be determined and recorded into a database, which can comprise a database text file or Excel spreadsheet, for example. However, embodiments can use any recording means without departing from the scope of the present invention. Values measured can be compared to the visual image. One or more types of numerical descriptors can be generated from the values. For example, descriptors such as a number of objects, an average, a standard deviation of objects, a histogram (number or percentage of objects per bin, average, standard deviation), and the like can be determined.

In a particular embodiment according to the present invention, data can be analyzed using morphometric values derived from any of a plurality of techniques commonly known in the art. For example, a software package called MetaMorph Imaging System, provided by Universal Imaging Corporation, a company with headquarters in West Chester, PA and NIH Image, provided by Scion Corporation, a company with headquarters in Frederick, Maryland.

30 Fluorescent images can be described by numerical values, such as for example, an area, a fluorescence intensity, a population count, a radial dispersion, a perimeter, a length, and the like. Further, other values can be derived from such measurements. For example, a shape factor can be derived according to a relationship

$4\pi * \text{area} / \text{perimeter}$. Other values can be used in various embodiments according to the present invention. Such values can be analyzed as average values and frequency distributions from a population of individual cells.

In a particular embodiment according to the present invention,
5 techniques for the automatic identification of mitotic cells are provided. Image analysis techniques employing techniques such as multidimensional representations, frequency-based representations, multidimensional cluster analysis techniques and the like can be included in various embodiments without departing from the scope of the present invention. Techniques for performing such analyses are known in the art and
10 include those embodied in MatLab software, produced by MathWorks, a company with headquarters in Natick, MA.

Scalar values providing efficacious descriptors of cell images can be identified using the techniques of the present invention to perform predictive analysis of drug behavior. In a presently preferred embodiment, a plurality of heterogeneous
15 scalar values can be combined to provide descriptors for each manipulation. By applying predictive analysis routines to the collections of these descriptors, predictive information about any number of manipulations and cell interactions can be extracted.

Fig. 7E illustrates a representative block flow diagram of simplified process steps for analyzing image feature values to obtain descriptors of cell state of
20 step 718 of Fig. 7B in a particular embodiment according to the present invention. This diagram is merely an illustration and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize other variations, modifications, and alternatives. Fig. 7E illustrates an input data of descriptors of known manipulations 319. A step 320 of reformatting and transforming data 319 to
25 formats suitable for analysis is performed. Additionally, a "cleaning" process can eliminate outlying data points and the like in the data. Then, in a step 322, a decision is made whether to continue with step 324 or with step 326 based upon determining a particular type of analysis appropriate for the present application or particular type of prediction. If decisional step 322 determines processing should continue with step
30 324, then, in that step, an error estimate using a set of test descriptors is performed to estimate the quality of a prediction and processing continues with step 320. Once an optimal prediction is achieved, processing continues with step 326. In step 326, optimal transformation parameters and prediction methods are selected for use in

steps 328 and 330 which analyze data about an unknown manipulation. In a step 328, a solution is generated based upon any of techniques including training a neural network, solving a mathematical equation, applying decision tree rules and/or the like. In a step 330, an input data set of unknown descriptors 318 is reformatted and
5 transformed based upon the optimal transformation parameters selected in step 326 using the transformation procedures in steps 320, 322 and 324. In a step 332, predictions techniques are applied to the reformatted manipulations from step 330 and the solution generated in step 328 and a plurality of properties of known manipulations 317 (e.g., therapeutic properties, and the like) in order to determine a
10 prediction of properties of unknown manipulation 316.

Fig. 7F illustrates a representative block flow diagram of simplified process steps for a method of mapping a manipulation of cells to a physiological characteristic in a particular embodiment according to the present invention. This diagram is merely an illustration and should not limit the scope of the claims herein.
15 One of ordinary skill in the art would recognize other variations, modifications, and alternatives. The method is generally outlined by the steps below:

(1) In a step 750, a plurality of cells, e.g., dead, live, cell fractions or mixtures of cells are provided.

(2) Then, in a step 752, the plurality of cells is manipulated, where
20 manipulation occurs using a source(s) from one or a combination selected from an electromagnetic, electrical, chemical, thermal, gravitational, nuclear, temporal, or a biological source.

(3) Next, in a step 754, a feature value is captured from the plurality of cells. The feature value can include one or any combination of characteristics such as
25 cell count, area, perimeter, length, breadth, fiber length, fiber breadth, shape factor, elliptical form factor, inner radius, outer radius, mean radius, equivalent radius, equivalent sphere volume, equivalent prolate volume, equivalent oblate volume, equivalent sphere surface area, average intensity, total intensity, and optical density. This list is not meant to be limiting.

(4) Then, in a step 756, a degree of presence of one or more feature values is assigned for each manipulation.
30

(5) In a step 758, the feature values from the plurality of cells are stored in memory locations. From the memory locations the values can be used for

statistical analyses to produce predictive information about the relatedness of the descriptors of the manipulations to one another. This information is used to infer properties of the manipulations.

Fig. 7G illustrates a representative block flow diagram of a simplified process steps for a method for populating a database with manipulated biological cell information in a particular embodiment according to the present invention. This diagram is merely an illustration and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize other variations, modifications, and alternatives. The method is generally outlined by the steps below:

(1) In a step 760, a plurality of cells in various stages of the cell cycle, A montage image that was used as a source to generate data in Appendix A is presented in Fig. 12., such as for example, the stages of interphase, prophase, metaphase, anaphase, and telophase are provided.

(2) Then, in a step 762, each of the cells in the various stages of mitotic development is manipulated.

(3) Next, in a step 764, an image of the plurality of manipulated cells is captured using image acquisition techniques in order to provide a morphometric characteristic of each of the manipulated cells.

(4) As a preferable option, in a step 766, an image database may be populated with the image of the plurality of manipulated cells.

(5) Following step 764 or optional step 766, a morphological value is calculated from the image in a step 768.

(6) In a step 770, the database is populated with the morphological value.

Fig. 7H illustrates a representative block flow diagram of simplified process steps for a method for populating a database with manipulated biological information, e.g., image acquisition parameters, image feature summary information, and well experimental parameters in a particular embodiment according to the present invention. This diagram is merely an illustration and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize other variations, modifications, and alternatives. Fig. 7H illustrates a step 780 in which cells are placed into site on a plate and a manipulation is applied. Then, in a step 781 an image is taken of the cells. In step 782, the image is transferred to an image archive

database. Then, in a step 783, well experimental parameters are entered into the database 787. Well experimental parameters can include cell type, manipulation and the like. In a step 784, image acquisition parameters are transferred to database 787. Image acquisition parameters can include file name, fluorophores and the like. In a
5 step 785, the image acquired in step 781 is analyzed. Then, in step 786, an image feature summary from the analysis step 785 is transferred to database 787.

In step 788, a lookup table for all analyses is provided to database 787. The lookup table provides information about the analyses. In a step 789, a query of database 787 for process data is performed. The results are reformatted. Then in a
10 step 790, the database 787 is queried. Next, in a step 791, features of the manipulations stored in the database are combined and reduced. Next, in a step 793, reduced features of step 791 can be compared. In a step 792, the results of step 793 are recorded in database 787. Then, in a step 794, a report of predictions based on comparisons performed in step 793 is generated.

15 Fig. 7I illustrates a representative block flow diagram of simplified process steps for acquiring images of manipulated biological information, e.g., cells, cell tissues, and cell substituents in a particular embodiment according to the present invention. This diagram is merely an illustration and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize other variations,
20 modifications, and alternatives. Fig. 7I illustrates a step 770 in which a user sets up an image analysis procedure. Then, in a step 772, an image is read into image analysis software. Next, in a step 774, patterns and objects are identified in the image using one or more algorithms. Next, in a step 776, sets of features are extracted from the image. Then, in a step 778, feature information, descriptor values and the like are
25 exported to the database, such as database 787 of Fig. 7H, for recording. Next, in a decisional step 779, a determination is made whether any more images should be taken. If this is so, processing continues with step 772. Otherwise, image acquisition processing is completed.

Fig. 7J illustrates a representative block flow diagram of simplified
30 process steps for populating, acquiring and analyzing images of manipulated biological information in a particular embodiment according to the present invention. This diagram is merely an illustration and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize other variations,

modifications, and alternatives. Fig. 7J illustrates a step 300 of placing a plate onto an imaging stage and reading a bar code. Then, in a step 301 an autofocus procedure is performed. Next, in a step 302, a first optical filter configuration is selected and an image is collected. Then, in a decisional step 303, a determination is made whether
5 more than one image per optical configuration can be taken. If so, then, in a step 304, a new position within the well is targeted and another image is collected. Then, in a decisional step 305, a determination is made whether any more images need to be collected. If this is so, step 304 is repeated until all images for a particular well have been collected. After one or more images are collected for the well, in a step 306, the
10 stage is returned to a starting position within the well, and a montage is created from collected images. The results are named with a unique file name and stored.

In a decisional step 307, a determination is made whether any more optical channels in the well can be imaged. If this is so, then in a step 308 the next optical filter configuration is selected and an image is collected. Processing then
15 continues with decisional step 303, as described above. Otherwise, if no further optical channels in the well can be imaged, then in a decisional step 309 a determination is made whether any wells remain to be imaged. If not all wells have been imaged, then in a step 310, the stage moves to the next well and processing continues with step 301, as described above. Otherwise, if all wells on the plate have
20 been imaged, then in a decisional step 311, a determination is made whether any more plates can be processed. If this is so, then processing continues with step 300 as described above. Otherwise, in a step 312, the information is stored to a CD or other storage device as a backup.

Fig. 7K illustrates a representative block flow diagram of simplified
25 process steps compound based upon information about effects of one or more known compounds on a cell population in a particular embodiment according to the present invention. This diagram is merely an illustration and should not limit the scope of the claims herein. One of ordinary skill in the art would recognize other variations, modifications, and alternatives. Fig. 7K illustrates a step 340 of populating a database
30 with descriptors for known compounds. Such descriptors can be determined from imaging the cell population. However, in some embodiments, descriptors can be derived by measurements and combinations of measurements and the like. Then, in a step 342, descriptors for the unknown compound are determined from imaging a

second cell population. The second cell population has been treated with the unknown compound. Then, in a step 344, a relationship between the descriptors determined from the unknown compound with the descriptors determined from the known compounds can be determined. Finally, in a step 346, an inference can be made about the unknown compound based upon the descriptors of the known compounds from the relationship determined in step 344.

Accordingly, the present invention provides a novel database design. In a particular embodiment according to the present invention, a method for providing a database comprises measurement of a potentially large number of features of one or more sub-cellular morphometric markers. Markers can be from any of a large variety of normal and transformed cell lines from sources such as for example, human beings, fungi, or other species. The markers can be chosen to cover many areas of cell biology, such as, for example markers comprising the cytoskeleton of a cell. The cytoskeleton is one of a plurality of components that determine a cell's architecture, or "cytoarchitecture". A cytoarchitecture comprises structures that can mediate most cellular processes, such as cell growth and division, for example. Because the cytoskeleton is a dynamic structure, it provides a constant indication of the processes occurring within the cell. The cytoarchitecture of a cell can be quantified to produce a one or more scalar values corresponding to many possible cellular markers, such as cytoskeleton, organelles, signaling molecules, adhesion molecules and the like. Such quantification can be performed in the presence and absence of drugs, peptides, proteins, anti-sense oligonucleotides, antibodies, genetic alterations and the like. Scalar values obtained from such quantification can provide information about the shape and metabolic state of the cell.

In a presently preferred embodiment, scalar values can comprise morphometric, frequency, multi-dimensional parameters and the like, extracted from one or more fluorescence images taken from a number of cellular markers from a population of cells. Two or more such scalar values extracted from a plurality of cell lines and markers grown in the same condition together comprise a unique "fingerprint" or descriptor that can be incorporated into a database. Such cellular descriptors will change in the presence of drugs, peptides, proteins, antisense oligonucleotides, antibodies or genetic alterations. Such changes can be sufficiently unique to permit a correlation to be drawn between similar descriptors. Such

correlations can predict similar properties or characteristics with regard to mechanism of action, toxicity, animal model effectiveness, clinical trial effectiveness, patient responses and the like. In a presently preferred embodiment, a database can be built from a plurality of such descriptors from different cell lines, cellular markers, and
5 compounds having known mechanisms of action (or structure, or gene response, or toxicity).

The present invention also provides database and descriptor comparisons according to other embodiments. In a particular embodiment according to the present invention, measurement of scalar values or features can provide
10 predictive information. A database can be provided having one or more "cellular fingerprints" comprised of descriptors of cell substance interactions of drugs having known mechanisms of action with cells. Such descriptors can be compared using a plurality of techniques, such as a technique of creating "phylogenetic trees" of a statistical similarity between the descriptors from various drugs. In a present
15 embodiment, scalar, numeric values can be converted into a nucleotide or amino acid letter. Once converted into a corresponding nucleotide representation, the descriptors can be analyzed and compared using software and algorithms known in the art for genetic and peptide sequence comparisons, such as GCG, a product of Genetics Computer Group, with company headquarters in Madison WI. In an alternative
20 embodiment, numeric values for the fingerprints can be used by comparison techniques. A phylogenetic tree can be created that illustrates a statistical significance of the similarity between descriptors for the drugs in the database. Because the drugs used to build the initial database are of known mechanism, it can be determined whether a particular scalar value in a descriptor is statistically predictive. Finally, a
25 compound fingerprint with no known mechanism of action can be queried against the database and be statistically compared and classified among the drugs in the database that the compound most resembles.

In a particular embodiment, relationships between measured morphometric properties and features of images and physiological conditions can be
30 determined. Relationships can include, for example, treatment of different cell lines with chemical compounds, or comparing cells from a patient with control cells, and the like. In a presently preferable embodiment, a clustering can be performed on acquired image descriptors. Some embodiments can comprise statistical and neural

network - based approaches to perform clustering and comparisons of various descriptors. The foregoing is provided as merely an example, and is not intended to limit the scope of the present invention. Other techniques can be included for different types of data. In some embodiments, clustering and comparing can be performed on features extracted from cell images. In a presently preferable embodiment, procedures for comparisons and phylogenetic analysis of biological sequences can be applied to data obtained from imaging cells.

Select embodiments comprising such approaches enable the use of a broad array of sophisticated algorithms to compare, analyze, and cluster gene and protein sequences. Many programs performing this task are known to those of ordinary skill in the art, such as for example, the program Phylip, available at <http://evolution.genetics.washington.edu/phylip.html>, and other packages listed at <http://evolution.genetics.washington.edu/phylip/software.html>. However, select embodiments according to the present invention can comprise a technique of statistical classification, statistical clustering, distance based clustering, linear and non-linear regression analysis, self-organizing networks, and rule-based classification.

Embodiments can perform such analysis based upon factors such as numerical value, statistical properties, relationships with other values, and the like. In a particular embodiment, numbers in a numerical descriptor can be substituted by one or more of nucleic acid or amino acid codes. Resulting "pseudo-sequences" can be subjected to analysis by a sequence comparison and clustering program.

Other types of databases can also be provided according to other embodiments. The database includes details about the properties of a plurality of standard drugs. When the descriptor of a test compound is compared to the database, predictions about the properties of the test compound can be made using any known property of the other compounds in the database. For example, properties about a compound in the database could include structure, mechanism of action, clinical side effects, toxicity, specificity, gene expression, affinity, pharmacokinetics, and the like. The descriptor of a compound of unknown structure from a natural products library could be compared to the descriptors of compounds with known structure and the structure could be deduced from such a comparison. Similarly, such information could lead to better approaches to drug discovery research including target validation

and compound analogizing, as well as pre-clinical animal modeling, clinical trial design, side effects, dose escalation, patient population and the like.

According to the present invention, databases can be integrated with and complementary to existing genomic databases. Differential genomic expression
5 strategies can be used for drug discovery using database technology. In one particular embodiment, cell data and cellular response data can be associated with a genetic expression profile assay to form a single assay. Live cells expressing fluorescence markers can be treated with a drug, imaged and analyzed for morphometry; and then analyzed for mRNA for expression. Such embodiments can provide rapid
10 development of tools to link cellular behavior with functional genomics.

Database methods according to the present invention can be used to predict gene function and to assist in target validation. Databases that include genetic diversity, i.e., having cellular descriptors from cells of differing genetic backgrounds (tumor, tissue specific, and gene knock out cell lines), can provide the capability to
15 compare cells of unknown genetic background to those in the database. Similarly, the descriptor of an unknown cellular portion in the presence of multiple drugs can be queried against the descriptors of the known markers in the database. For example, if an unknown gene is tagged with Green Fluorescent Protein (GFP), the database may be used to identify the cellular portions for which that unknown gene encodes.

According to the present invention, target validation and specialized
20 cell-based assay screening can be performed using database systems and methods to serve as a universal high-throughput cell-based assay that can evaluate the molecular mechanism of drug action. As new genes are isolated and identified, a large collection of available gene-based knowledge is becoming available. From this large
25 collection of new genes, potential protein targets can be identified using the genomic tools of sequence analysis and expression profiling. However, unless a gene mutation is tightly linked to a disease state, further validation of individual targets is a time consuming process, becoming a bottleneck in drug discovery. Furthermore, robotics and miniaturization are making "High Throughput Screening (HTS)" the industry
30 standard, substantially reducing the time and cost of running a target-based biochemical assay. Therefore, it is now possible to routinely screen large libraries and use a resulting "hit" to validate the target. In such approaches, a specialized cell-based assay would be developed to test hits for each target. Since this often involves

the creation of cell lines expressing new markers, this stage may also become a bottleneck that cannot keep pace with HTS. In addition, these cell-based assays may not be amenable to high-throughput screening, making it difficult to test the increasing number of analogs arising from combinatorial chemistry.

5 In a particular embodiment according to the invention, a rapid characterization of large compound libraries for potential use as pharmaceutical products can be provided by predicting properties of compounds that relate to the compounds' potential as bioactive drugs. In many drug discovery situations, virtually millions of compounds can be passed through a HTS assay against a small number of
10 validated targets. These assays produce hundreds to thousands of potential hits. These hits can then be subsequently screened by a pipeline of secondary and tertiary screens to further characterize their specificity, often time completely missing non-specific interactions with other proteins. Techniques according to the present invention can provide a replacement to such screening operations by providing
15 information about cellular accessibility and mechanism of action for the hits coming from a HTS system. Furthermore, it can replace the biochemical HTS assay and allow rapid and accurate identification of attractive compounds from large libraries without an intervening biochemical assay. The cell information can be predictive of whether to continue into an animal model for each compound, and which animal model to
20 pursue.

 The principles of the present specifically contemplate a wide variety of research methodologies, or usage scenarios, implementing these principles. The following discussion of three such scenarios is by way of illustration and not limitation. Study of the principles enumerated herein will render evident to those
25 skilled in the art certain additional methodologies or usage scenarios enabled by the teachings hereof. The present invention specifically contemplates all such modifications. The following description presents some specific embodiments and scenarios that represent a broader use of cellular phenotypic data and characterizations to deduce mechanisms of action and other features of cellular
30 responses to various stimuli. Such procedures generally involve producing a quantitative cellular phenotype based upon two or more cellular attributes and then comparing that phenotype to phenotypes previously stored and indexed. Such

procedures make use of databases or other repositories of biological information. The invention is not limited to the specific embodiments described here.

Considering first the procedure 2000 depicted in Figure 20, a compound has been identified as having a particular cellular activity. See 2004. For example, a compound may be found to inhibit the growth of certain cancer cell *in vitro* by a specific and desired mechanism of action. This may be a particular company's "gold standard."

Next, the compound is analyzed at 2006 in terms of its effect on one or more cell lines. More specifically, the compound is linked, virtually, to a particular phenotype. Two or more values or measures of cellular attributes characterize that phenotype. These attributes are quantified in the context of specific cellular markers.

In one example, the cellular marker is an organelle such as a nucleus or Golgi apparatus. Measured attributes useful for characterizing an associated phenotype include geometric parameters (e.g., size, shape, and/or location of the organelle) and composition (e.g., concentration of particular biomolecules within the organelle).

The phenotype may be characterized by administering the compound of interest to various cell lines and in various concentrations. In each example within this matrix, the attributes of interest are measured. Ultimately, certain phenotypic features (combinations of attribute values) are associated with the compound of interest. These features provide a template for the phenotype.

Next, using the phenotype as identified at 2006, the process identifies other compounds providing similar features. The goal here is to present a list of compounds having a mechanism of action similar to that of the compound that started the process. This allows researchers to identify a mechanism of action, if not already known, for their compound and to draw conclusions based upon their compound's link to other known compounds (which may not be chemically/structurally similar to the compound of interest).

Identifying similar compounds based upon phenotype can take many paths. Most will involve some mathematical basis. For example, the phenotype defined at 2006 can be represented as a fingerprint or vector comprised of multiple scalar values of cellular attributes (as described above). The phenotype representation can then be compared against known phenotypes characterized by the same format

(e.g., they are all characterized as vectors having the same attribute set, but with different values of the attributes). The comparison may be as simple as a Euclidean distance or more sophisticated as a neural network or multivariate statistical correlation.

5 The known compounds and associated phenotypes may be stored as database records or other data structures that can be queried or otherwise accessed as part of the identification procedure. The compounds may also be associated with other relevant data such as clinical toxicity, cellular toxicity, hypersensitivity, mechanism of action, etc. (when available).

10 Compounds found to be sufficiently similar to the starting compound are returned for consideration by researchers. A data processing system may rank such compounds based on degree of similarity to the starting compound. In some cases, the system may even provide similarity scores associated with the listed compounds.

15 Often researchers wish to determine whether their particular compound has clinical or biochemical effects beyond those that they are already aware of. In a typical scenario, the compound of interest was selected based upon its strong binding a target or its stimulation or inhibition of cell growth in a particular cell line. The process associated with 2010 has likely identified the compound of interest as having
20 a particular mechanism of action based on phenotypic similarity to other compounds having a similar mechanism of action. However, within the region of biochemical space, there may be subspaces (characterized by subphenotypes) that correspond to separate properties. For example, within the phenotypic space associated with one mechanism of action, there may be subspaces associated with clinical toxicity,
25 cellular toxicity (likely overlapping the clinical toxicity space), and little or no toxicity. Obviously, a researcher would like to know whether her compound is likely to be toxic.

 Thus, the process 2000 may include characterizing the compound of interest in terms of its distance from (i.e., similarity to) specific phenotypes having
30 known characteristics. In a typical example, the known characteristic is toxicity. This feature allows the researcher to quantify her compound in terms of mechanism of action AND toxicity (or in terms of two or more other relevant properties associated

with phenotype). To allow simple ranking or characterization, compounds of interest may be scored according to a simple or weighted Boolean expression.

A second scenario of interest is depicted in Figure 21. This scenario again defines a phenotype in terms of a quantifiable vector or other measure.

- 5 However, rather than using a compound of interest to generate the phenotype, some other cellular stimulus is used to generate the phenotype.

As shown, a process 2100 begins with receipt of cells of interest. See 2104. In many situations, the cells are produced by a genetic or epigenetic process that affects the expression level or activity of a particular protein. More generally,
10 any cellular stimulus (e.g., radiation level and type, gravity level, magnetic field, acoustic perturbations, etc.) can be used to generate the cell line of interest. Importantly, this stimulus affects the phenotype and can be correlated therewith.

In the context of drug discovery, a gene encoding for a particular target can be genetically knocked out, underexpressed, overexpressed, expressed in a non-
15 native state, etc. This may be accomplished via standard procedures involving genomic modification, translation or transcription apparatus modification (e.g., use of antisense nucleic acids), blocking target activity (using antibodies to a receptor site for example), and the like. These processes will generally affect the phenotype in some quantifiable way. Importantly, they clearly and unambiguously define a cellular
20 phenotype associated with altering the activity of the target protein.

At 2106, the process involves measuring one or more cellular features from the cell line of interest to define/quantify the phenotype. This may be accomplished as described above with reference to 2006. Next, at 2108, the cellular phenotype generated in this manner is used to identify and rank a set of compounds
25 associated with the phenotype. This operation may proceed in the manner of operations 2008 and/or 2010 from Figure 20.

Finally, at 2110, the process clusters the compounds returned at 2108 by a mechanism of action. The operation 2106 has tightly bound a mechanism of action to a phenotype. Various compounds characterized and stored in a system
30 database may be tentatively assigned a mechanism of action or may have no suggested mechanism of action. By matching their virtual phenotype to the phenotype generated at 2106, one can create or strengthen an association between the compounds and mechanism of action relevant to the stimulus at 2104.

Considering now Figure 22, a third scenario is depicted. This scenario again involves using a virtual phenotype to glean information relevant to a mechanism of action or other cellular activity. In this case, assay data from a group of compounds (e.g., a primary or focused library) is used to elucidate a phenotype.

5 As shown, a process 2200 begins by identifying a target protein. See 2204. Then, at 2206, the process involves identifying positive and negative biochemical hits. More generally, this may involve ranking a number of compounds based upon their interaction with the target. In a specific case, the compounds are ranked based upon their binding affinities to or ability to inhibit the enzymatic activity
10 of the target protein.

 After the compounds have been characterized in some manner based upon their interaction with the target, they are used to define a cellular phenotype. See 2208. Generally, the techniques to accomplish are the same as described with reference to operation 2006 of Figure 20. In this case however, one may obtain a
15 strong correlation between mechanism of action (involving the target) and phenotype by using multiple of the compounds identified at 2206. For example, some of the “best hits” may be administered to cell lines in various concentrations. And some of the least effective compounds may also be administered. Cellular attributes that are more strongly exhibited with increasing concentration of the best hits (and not
20 exhibited or exhibited only weakly upon administration of the negative hits) can be used to define the virtual phenotype. In a related approach, compounds having widely varying levels interaction with the target are administered to cells. Those cellular attributes that vary linearly or at least monotonically with the degree of interaction between the target and compound represent attributes that can be used to define the
25 virtual phenotype.

 After the cellular phenotype has been defined, previously characterized compounds may be clustered with that phenotype. See 2210. As with operation 2110 of Figure 2, this may create or strengthen an association between a mechanism of action and various compounds in a database.

30 Finally, and optionally, procedure 2200 may provide a “higher resolution” mechanism of action for the compounds identified at 2206. See 2212. Presumably interaction with the target suggests a specific mechanism of action or at least some aspect of a mechanism of action. However, a given target may participate

in a larger cellular mechanism of action – unknown to researchers. Further, a compound may that binds with the target may participate in multiple mechanisms of action – some of which do not involve the target. By linking the target (and its positive hits) to a particular phenotype, some of these additional cellular level activities can be elucidated. The defined phenotype may have been previously identified as associated with other mechanisms of action or higher resolution mechanisms of action. Thus, the phenotype identified at 2208 can be leveraged to generate a higher resolution mechanism of action at 2212.

As suggested in the above discussion, compounds and associated phenotypes may be stored as database records. Such databases can take on many flavors. In one example, a database includes various pieces of information relevant to oncology. Such database may include numerous compounds classified by cellular phenotype, mechanism of action, toxicity, etc. More specifically, the database may include data on commercially available compounds clustered by cellular phenotypes corresponding to mechanisms of action. Further the databases of interest may extended or combined (via standard relational tables and algebra for example) to include additional data such as pharmacology data, cellular genomics data, gene expression data, protein expression data, etc. In a specific example, the database includes measurements made on a subset of the NCI60 cell lines, using DNA, Golgi apparatus, and/or microtubules as markers for defining the phenotypes. Other data includes dosage response information, variation in effect over time, etc. The compounds populating the database could include known National Cancer Institute oncology study compounds. In a specific embodiment, the compound set includes some or all of the compounds mentioned in the article “A gene expression database for the molecular pharmacology of cancer,” Nature Genetics, 24, pp. 236-244 (March 2000).

Various biological analyses may be conducted to develop additional information for characterizing compound mechanisms of action, etc. For example, a cell count analysis may be used to develop dose response curves, GI 50 data, etc. The cell cycle may also be analyzed to find out how various stages in the cycle vary in response to particular stimuli. The Golgi apparatus may be analyzed to determine whether it is in a normal state, a dispersed state, a diffused state, etc. As another example, tubulin may be analyzed to determine whether it is normal, de-polymerized,

over-polymerized, bundled, etc. Obviously, combinations of such analyses may be performed. For example, properties of the Golgi apparatus or tubulin may be analyzed over one or more cell cycles.

In some embodiments, techniques according to the present invention
5 can provide tools for the later stages of drug development such as clinical trial design and patient management. The properties of known drugs, such as clinical trial and patient response information, will be used in a similar fashion as the pre-clinical information to provide predictions about the properties of novel compounds. Because the human cell is the locus of drug action, a database containing drug-cell interactions
10 will be able to provide predictive value for this aspect of drug development.

Although the above has generally been described in terms of specific hardware, software, and methods, it is understood that many alternatives can exist. In particular, the present invention is not limited to a particular kind of data about a cell, but can be applied to virtually any cellular data where an understanding about the
15 workings of the cell is desired. Thus, in some embodiments, the techniques of the present invention could provide information about many different types or groups of cells, substances, and genetic processes of all kinds. Of course, one of ordinary skill in the art would recognize other variations, modifications, and alternatives. Some examples according to the present invention are provided below.

20

EXPERIMENTS

To prove the principle and demonstrate the objects of the present invention, experiments have been performed to determine the effects of manipulations on cell structure using imaging and analysis techniques applied to a variety of
25 situations. These experiments were performed by growing multiple cell lines in the presence of multiple compounds, or substances. Cells were fixed and stained with fluorescent antibodies or labels to multiple cellular portions. One or more images of the cells were then obtained using a digital camera. Descriptors were built by quantifying and/or qualifying patterns of one or more feature from each image in the
30 cell lines under study. A database was built from the descriptors. As the database grows, it should be able to predict the mechanism of action of an unknown drug by comparing its effect with the effects of known compounds or to identify data clusters within large libraries of compounds.

In a first experiment, an automated method to count the number of cells and differentiate normal, mitotic, and apoptotic cells was created.

Approximately, 5,000 HeLa cells were plated per well in a 96 well plate and grown for 3.5 days. The cells were fixed with -20° MEOH for 5 minutes, washed with TBS
5 for 15 minutes, and then incubated in 5 mg/ml Hoechst 33342 in TBS for 15 minutes. Then, 72 images were collected with a 40x objective and 75 ms exposure time.

The analysis was performed on objects that met a certain size criteria that was based on 1) measuring the size of objects in the image that were clearly not cells and 2) excluding the first peak of the area histogram (Fig. 8B values 1-4654).

10 Histograms of the individual object data were generated for each type of feature. Fig. 8A shows the histogram for average intensity, and Fig. 8B shows histogram data for the area of each object. Fig. 8C shows the scatter plot of the average intensity vs. the area of all of the objects. The pattern of the scatter plot showed an interesting pattern: a large cluster of cells in one region of the graph, with
15 a scattering of object points in other regions. Because mitotic structures are identified as particularly bright objects, most likely due to the biological fact that the chromatin is condensed, the original Hoechst images could be used to identify which cells were either undergoing mitosis, or otherwise looked abnormal. Manual inspection of 917 cells resulted in the classification of each object. Fig. 8D shows a graph where each
20 type of cellular classification is delimited. This graph clearly shows that the mitotic nuclei are brighter than the interphase nuclei. Further, the different phases of the cell cycle can be separated using these two features. Figs. 8E-8F show bar graphs of the average and standard deviations of the areas and average intensities for each cell classification type. These graphs show that interphase nuclei are statistically less
25 bright than mitotic nuclei and that telophase nuclei are statistically smaller than other mitotic nuclei.

Each image was thresholded to an intensity level of 20. A standard area value was set at 9500 pixels. Automated information gathering about all of the objects was done and collected into an Excel spreadsheet (for more information see,
30 section on imaging system). The following information was recorded:

IMAGE NAME
OBJECT #

AREA
STANDARD AREA COUNT
PERIMETER
FIBER LENGTH
FIBER BREADTH
SHAPE FACTOR
ELL. FORM FACTOR
INNER RADIUS
OUTER RADIUS
MEAN RADIUS
AVERAGE INTENSITY
TOTAL INTENSITY
OPTICAL DENSITY
RADIAL DISPERSION
TEXTURE DIFFERENCE MOMENT
EFA HARMONIC 2, SEMI-MAJOR AXIS
EFA HARMONIC 2, SEMI-MINOR AXIS
EFA HARMONIC 2, SEMI-MAJOR AXIS
ANGLE
EFA HARMONIC 2, ELLIPSE AREA
EFA HARMONIC 2, AXIAL RATIO
EFA HARMONIC 3, SEMI-MINOR AXIS

The following results were obtained:

- 1,250 objects were counted
- 201 of those objects has standard area counts > 2 (area > 19000 pixels)
- 195 objects had areas < 6000 pixels
- 1529 objects estimated in total
- 1328 object areas are > 6000 pixels
- The data was reduced to 917 objects that were $6000 < \text{area} < 19000$
- For the 917 objects a scatter plot of area vs. average intensity and a histogram of the average intensity were generated.

- 116 objects that had average intensity intensities > 60 were manually looked at to determine their morphology.
 - Of those 116 objects:
 - 6 were dead or indistinguishable
 - 4 were interphase
 - 30 were prophase
 - 32 were metaphase
 - 24 were anaphase
 - 20 were telophase (10 pairs)
- 10
- 12 prophase objects were missed because of gray scale cut off. (8 of those prophase cells had gray scale values > 57 , as did 7 interphase)
 - 1 telophase object was missed because it was too small (< 6000)
 - 1 prophase object was missed because it was too big (> 1900)
- 15
- 16 mitotic objects were missed because they were parts of objects with standard count > 2 .

In sum, out of 917 single objects, the analysis correctly identified 106 out of 130 mitotic objects, or (81% predictive, 91% of identified mitotics). Out of 917 single objects, the analysis incorrectly identified only 10 non-mitotics as mitotics (1% total, 8% of identified mitotics); 14 mitotics as interphase (1.4% total, 1% interphase). An automated classification system that would automatically assign values to each object using these or other measurement features can thus be developed, utilizing the principles set forth herein.

In a second experiment, the effects of Taxol on MDCK cells and the different types of morphological effects were observed. A plurality of MDCK cells grown in 96 well plates were treated with Taxol for 4.5 hours at different concentrations (10 uM-1pM). They were then fixed, labeled with Hoechst, and imaged.

This experiment used a labeling protocol comprising: MEOH fix at – 20°, Wash in PBS, Block in PBS/BSA/Serum/Triton-X 100, Incubate with 5 µg/ml Hoechst 10 minutes, and wash.

Cells were inspected for different morphologies and manually counted at each different drug concentration in one well. Fig. 9 shows example images from each drug concentration and the different types of morphologies and cells are highlighted. Fig. 10 shows the distribution of each morphology within the cell population as a function of drug concentration. The higher the concentration of Taxol, the larger proportion of cells underwent apoptosis, and the fewer number of normal mitotic cells were detected.

In a third experiment, the purpose was to determine whether the automated analysis methods developed in the first experiment can detect differences in Hoechst morphology in the presence of 6 known compounds at one concentration and exposure time in one cell line. In this experiment, HeLa cells were separately treated with 6 compounds with known mechanism of action. The quantitative methods described in the first experiment were applied to the Hoechst images.

Approximately 5,000 HeLa cells per well were plated in a Costar black-walled 96 well tissue culture treated plate and left to recover in the incubator for 24 hours. After this time, 10 ug/mL of cytochalasin D (CD), Taxol, hydroxyurea, vinblastine, nocodazole, and staurosporine was added to different wells at a 1:100 addition in DMSO.

The cells were incubated in the presence of drug for 24 more hours. After 24 hours, the cells were removed and fixed as in the first experiment. Then, 9 images per well were collected of the Hoechst staining using a 10x objective.

The low magnification images taken of Hoechst were run through the automated image analysis method described in the first experiment. Plots of the average intensity and area were made of each compound. Fig. 11 shows the scatter plots of the compounds. The scatter plots of each compound are visually distinct. For example, cells treated with CD are smaller than control, and cells treated with Hydroxyurea are larger and brighter. Furthermore, the number of cells per well was very different (data not shown).

The effects of different compounds can be clearly and automatically distinguished by identifying changes in cellular morphology. This method can also be used to count adherent cells.

The next experiment was to develop clustering algorithms that assign statistically meaningful values to the representative two dimensional data shown in Fig. 10, and even more complicated clustering of all of the multidimensional data that can be extracted across one, and multiple images.

A fourth experiment was performed to obtain high magnification images of two markers in the presence of drugs. In this experiment, HeLa cells were treated with 80 generic compounds with known mechanism of action. The quantitative methods described in the first experiment were applied to the Hoechst images.

Approximately 5,000 HeLa cells per well were plated in a Costar black walled 96 well tissue culture-treated plate and left to recover in the incubator for 24 hours. After this time, 10 ug/mL of each compound from the Killer Plate from Microsource Discovery Systems (Gaylordsville, CT) was added to different wells at a 1:100 addition in DMSO. The cells were incubated in the presence of drug for 24 more hours. After 24 hours, the cells were removed and fixed as in the first experiment. In addition to being labeled with Hoechst 33342 (against chromatin), cells were also labeled with 1 unit of rhodamine-conjugated phalloidin (against actin) for 30 minutes.

The 96 well plate was imaged twice. Once, 9 images per well were collected of the Hoechst staining using a 10x objective. After this, one image per well of both the phalloidin and Hoechst staining was collected using a 40x objective.

The resulting high magnification images were analyzed qualitatively and distinct pattern differences were detected in both the Hoechst and phalloidin images. Fig. 12 shows three example images from the experiment. The top row is the Hoechst staining, and the bottom row is the phalloidin staining from the same well. The columns show the images from wells treated with just DMSO (control), cytochalasin D, and Colchicine. The morphology of each marker is different in the presence of each drug. Interestingly, there is an effect in the morphology of the chromatin in the Hoechst image of cytochalasin D, which directly targets the actin cytoskeleton (and thus there is an expected effect in the phalloidin image). Also, there is an effect on the actin cytoskeleton, compared to control, in the presence of colchicine that directly targets the microtubule network.

The low magnification images were analyzed as described in the first experiment, and different patterns were seen in both the average intensity vs. area plots, and in the number of cells per well (data not shown). Thus, changes in patterns of a marker that is "down-stream" from the direct target of a compound are detectable. Automated image analysis protocols for actin and other markers can be developed similarly, again utilizing the principles set forth herein.

A fifth experiment was performed to test quadruple labeling of 9 different cell lines grown in normal conditions. In this experiment, NCI-H460, A549, MDA-MD-231, MCF-7, SK-OV-3, OVCAR-3, A498, U-2 OS, and HeLa cells were plated. Then, the cells were fixed and stained for portions of the each cell known as DNA, tubulin, actin, and Golgi.

The following table summarizes the procedures for this experiment:

Action	Active Ingredient/Notes	Buffer	Vol/ well	Desired Time	Temp
Remove media	NOTE: gently by pipetting, not aspiration				
Fix	4% Formaldehyde	PBS	100µl	20 min	rt
Wash		TBS	100µl	5 min	rt
Wash		TBS	100µl	5 min	rt

Permeablize	0.1% Triton X-100	TBS	100 μ l	10 min	rt
Permeablize	0.1% Triton X-100	TBS	100 μ l	10 min	rt
Block	% BSA % Serum Filter sterilize before use	TBS w/azide	100 μ l	1hr or o/n	rt or 4°C
Primary Antibody	1:1000 dilution of DM1 α	TBS + 1% BSA + 0.1% TX-100	50 μ l	1hr or o/n	rt or 4°C
Wash		TBS	100 μ l	5 min	rt
Wash		TBS	100 μ l	5 min	rt
Wash		TBS	100 μ l	5 min	rt
Fluorescent Stain	FITC lens culinaris 1:500 Rhodamine-Phalloidin 1:500 CY5 goat anti-mouse 1:100	TBS + 1% BSA + 0.1% TX-100	50 μ l	1 hr.	rt, dark
Wash		PBS	100 μ l	5 min	rt, dark
Hoechst	1:1000 dilution of 5mg/ml	TBS	100 μ l	15 min	rt, dark
Wash		PBS	100 μ l	5 min	rt, dark
Wash		PBS	100 μ l	5 min	rt, dark
Wash		PBS	100 μ l	5 min	rt, dark
Store		PBS	200 μ l	1 month	4°C

Cells were plated out at different densities for 48 hours. Cells were fixed and labeled by the above method. Cells were imaged using an automated imaging system that collected 9 images from each marker using a 10x objective.

Higher magnification images were collected of a few cells for demonstration purposes.

In this experiment, each cell line demonstrated different morphological patterns as determined by phase. For example, A549 cells are much more compacted than OVCAR-3 cells as determined by phase contrast imaging (data not shown). The different fluorescent markers showed even bigger differences between different cell lines. Figs. 13 and 14 show 4 panels of each marker for A549 (Fig. 13) and OVCAR-3 cells (Fig. 14). The markers are Hoechst (upper left), Phalloidin (upper right), Lens culinaris (lower left), and DM1a antibody (lower right). The following table summarizes the qualitative differences between these images:

MARKER	A549	OVCAR3
Hoechst/DNA	small	large
Phalloidin/actin	fuzzy	crisp - many stress fibers
Lens culinaris/Golgi	compact	Disperse/punctate
DM1alpha/Tubulin	perinuclear	evenly distributed

Higher magnification images were taken of the OVCAR3 cells. Fig. 15 shows the same markers at 20x, and Fig. 16 shows the markers at 40x. While the highest magnification images show the most detail, these images illustrate that very little morphological or feature information is lost in the 10x images.

These data exemplify the differences in morphology seen between different cell types. Thus the automated image analysis software can be customized for each marker in each cell type. Different drugs should effect these morphologies differentially.

An automated quantification method for each marker and cell line can be similarly developed.

A sixth experiment was conducted with a more sophisticated software package and to develop more flexible image recognition algorithms. In this experiment, prototype image features extraction was performed using MatLab programming language with image toolbox and SDC morphology toolboxes. Algorithms are being developed that will automatically identify objects on images and

to measure various morphological and feature parameters of these objects. Many different features for each of the cellular markers were acquired.

An example of a MatLab program called "AnalyseDNA" that takes as an input an unlimited number of images, identifies individual objects in these images based on either their intensities, or based on edge-detection algorithms, and extracts a number of morphological and intensity characteristics of these objects. A copy of this program follows:

Listing of the AnalyseDNA.m program and of some of the supporting subroutines

```
10 function files_analysed = AnalyseDNA(filemask, outpath,
    nx, ny, filter_range, dext, modifier, sfname) .
    % AnalyseDNA performs measurements on files of DNA images
    % V1. EV 2-11-99; 2-15-99; 2-16-99
15 %
    % files_analysed = AnalyseDNA(filemask, outpath, nx, ny,
    filter_range, dext, modifier, sfname)
    %
    % PARAMETERS:
20 %     ALL PARAMETERS ARE OPTIONAL
    %
    %     FILEMASK - mask for file names to be analyzed
    INCLUDING PATH(for example c:\images\*.tif)
    %     DEFAULT '.\*.tif' (all *.tif files in the current
25 directory) .
    %
    %     OUTPATH - path to a directory where all the output
    files will be placed.
    %     DEFAULT - output is saved in the same directory
30 which contains images
    %
    %     NX, NY - number of individual images in montage
    images along X and Y axes (DEFAULT 1)
    %
```

```
%    FILTER_RANGE - 3 col-wide array (or[]). Specifies
how data is filtered when summary is calculated
%    this parameter internally is passed to GetDNADData
and then to GetSummaryData - see these
5 %    functions for details. For example: [2 2 Inf; 6 100
8000] will case all raws of data for which
%    values in column 2 are less than 2 and all raws
where values in column 6 are less than 100 or
%    more than 8000 to be excluded from all
10 calculations of a summary.
%    DEFAULT - [] (means do not filter, summarize all
data)
%
%    DEXT - string. Extension for data files being saved.
15 %    DEFAULT 'dat';
%
%    MODIFIER - this modifier is 'SUMMARY', summary file
is created;
%    'SUMMARY ONLY' - only summary is generated,
20 data for individual files are not saved
%
%    sfname - string. File name of a summary file
%    DEFAULT 'summary[date].dat'
%
25 % OUTPUT:
%
%    AnalyseDNA works on image files or montages. For
each image file it creates a tab-delimits file of
measured
30 %    parameters of all the objects in the montage with
the same base name as a montage file and extension
specified
```

```
%    by dext parameter (or .dat by default) and file
'errors[date].err' - with the list of files that matched
the
%    filemask but could not be processed.
5 %    If 'summary' or 'summary only' modifier is
specified, it also creates a single file
'summary[date].dat' (or
%    different extension, if specified by DEXT) which
contains summary information for all analyzed files.
10 %
%    ALL OUTPUT FILES are saved in a directory specified
by OUTPATH parameter
%
%    RETURNS *files_analysed* - number of files that have
15 been successfully processed.
%
%    Column designations in the output files are
described in GetDNADData
%
20 % FILE NAME CONVENTIONS
%    AnalyseDNA attempts to identify a number for each
file to identify the file in summary output.
%    It does that by looking for the first space or
underscore, followed by a number and then takes
25 %    as many successive numbers as it can find. If it
fails to identify a number it assigns a
%    default which is -1
%
%
30 % SEE ALSO GetDNADData, GetSummaryData
%
% TO DO    improve error handling in opening and writing
files (GLOBAL error_file ?)
```

```
%           include procedures for writing text headers
into the output files

if nargin > 8
5   error ('Wrong number of input parameters');
end
if nargin > 1
    error ('Wrong number of output parameters: only one
allowed');
10 end

% set defaults
need_summary = 0;
summary_only = 0;
15 use_default_outpath = 0;
datestring = datestr(floor(now));
if nargin == 7      % set default summary file name
    sfname = ['summary' deblank(datestring)]; % extension
will be appended later based on dext
20   if deblank(upper(modifier)) == 'SUMMARY'
        need_summary = 1;
    elseif deblank(upper(modifier)) == 'SUMMARY ONLY'
        need_summary = 1;
        summary_only = 1;
25   else
        error (['Wrong parameter: unknown modifier '
modifier]);
    end
end
30
if nargin == 5
    % default data file extension
    set_dext = 'dat';
end
```

```
    if nargin == 4
        % default filter range
        filter_range = [];
    end
5   if nargin == 3
        ny = 1; % default number of images in montage along Y
    end
    if nargin == 2
        nx = 1;
10  end
    if nargin == 1
        use_default_outpath = 1;
    end
    if nargin == 0
15  filemask = '*.tif'
    end

    % check parameters
    if ( ~ischar(filemask) | ~ischar(dext) | ~ischar(sfname)
20  )
        error('Wrong parameter type: filename, filepath,
dext and sfname should be strings');
    end
    if ( ( size(nx) ~= [1 1] ) | ( size(ny) ~= [1 1] ) )
25  error ('Wrong parameter type: nx and ny should be
scalars (1x1 arrays)');
    end
    if (~isempty(filter_range) & size(filter_range, 2) ~= 3)
        error ('Wrong parameter type: filter range should be
30  [] or 3 - cols-wide array');
    end
    % end testing parameters

    % Generate list of files to process
```

```
datapath = getpath(filemask);
if use_default_outpath == 1
    outpath = datapath;
5  end
if exist(outpath, 'dir') ~= 7
    error(['Path ' outpath, 'not found. Exiting..']);
elseif exist(datapath, 'dir') ~= 7
    error(['Path ' datapath, 'not found. Exiting..']);
10 end

sfname = makefullname(outpath, sfname, dext);
if need_summary == 1
    if exist(sfname, 'file')
15     disp(['File ', sfname, 'already exists!']);
        input ('Press ^C to abort, Enter to delete and
continue');
        delete(sfname);
    end
20 end

flist = FileList(getfname(filemask), datapath);
numfiles = size(flist, 1); % total number of files to
25 process
disp(['About to process ', num2str(numfiles), ' files']);
%DEBUG - commented out "input" to run from Wrod
input('Press ^C to abort, Enter to continue');

30 % main loop where the job gets done:
error_file = makefullname(outpath, ['error' datestring
'.err']);
num_processed = 0;
num_error = 0;
```

```
for i = 1:numfiles
    % first generate file name for a data output file
    current_fullname = flist(i, :); % full name with path
    and extension
5    current_datafile = makefullname(outpath,
    makefname(getbasefname(current_fullname), dext) );

    %extract number from a filename
    fnumber = getfilenumber(current_fullname);
10
    % load an imagefile, record errors
    read_error = 0;
    try
        I = imread(current_fullname);
15        %DEBUG
        disp(['Image file #', num2str(fnumber), '
loaded']);
    catch
        % record file-opening error in an error_file
20        read_error = 1;
        num_error = num_error +1;
        msg = [current_fullname ': ' lasterrr];
        add_error_msg(error_file, msg);
    end
25
    % extract and write data to a file in outpath
    if read_error ~=1
        if (need_summary == 0)
            %DEBUG
30            disp(['Starting analysis of file #',
num2str(fnumber), '.']);
            current_data = GetDNADData(I, nx, ny, fnumber);
            %DEBUG
```

```

        disp(['Finished analysis of file #',
num2str(fnumber), '.']);
        %load current_data.mat 'current_data';
        write_data(current_data, current_datafile);
5      else      %summary needed
        %DEBUG
        [current_data, current_summary] = GetDNADData(I,
nx, ny, fnumber, filter_range);
        %load current_data.mat 'current_data';
10      %load current_summary.mat 'current_summary';
        write_summary(current_summary, sfname);
        if summary_only ~= 1
            write_data(current_data, current_datafile);
        end
15      end
    end
end % of the main for loop
num_processed = numfiles - num_error;

20 %=====end function AnalyseDNA()
=====

%=====
=====

25 function result = add_error_msg(filename, msg)
% adds string MSG to an errorfile FILENAME
% returns 1 if success, 0 if failure

err_FID = fopen(filename, 'at');
30 if err_FID == -1
    warning(['Can not open error file ' filename]);
else
    fprintf(err_FID, '%s\n', msg);
    fclose(err_FID);

```



```

end

%=====end function add_error_masg()
=====

5  %=====
=====

function N = getfilenumber(fname)
% returns the first number extracted from a file name
(string) or -1 if fails to extract any number
10 numbers = NumbersFromString( getfname(fname) ); % vector
of all numbers encoded in the name

                                     % (but not in the path, even if
present)
15 if isempty(numbers)
    N = (-1);    % return -1 if no numbers found in the
name
else
    N = numbers(1);
20 end

%===== end function getfilenumber()
=====

25 %=====
=====

function result = write_data(data_array, file_name)
% writes data in a data_array in a tab-delimited ascii
file.
30 % result is 0 if success and -1 if failure
% if file_name exists, overwrites it
result = -1;
try
    fid = fopen(file_name, 'wt');

```

```

        if fid ~= -1
            for k = 1:size(data_array, 1)
                fprintf(fid, '%g\t', data_array(k, :));
                fprintf (fid, '\n');
5         end
            test = fclose(fid);
            result = -1;
        catch
            result = -1;
10    end

%===== end function write_data()
%=====

15 %=====
%=====
function result = write_summary (s_vector, file_name)
% appends summary vector s_vector to a file_name (ASCII
tab-delimited file).
20 % if file_name does not exist, creates it.
% result is 0 if success and -1 if failure
%
result = -1;
try
25     % debug
        fid = fopen(file_name, 'at');
        result = fprintf(fid, '%g\t', s_vector);
        result = fprintf(fid, '\n');
        result = fclose(fid);
30     result = 0;
    catch
        result = -1;
    end

```

```

% ===== end function write_summary()
=====

function Data = GetObjectsData(I, Ilabel)
5 % GetObjectsData returns array measurements of objects in
  image "I" masked by "Ilabel"
  % EV 2-3-99; 2-10-99
  % OData = GetObjectsData(I, Ilabel) returns an array of
  morphological and intensity measurements
10 %   taken from a grayscale image "I". Objects are
  identified on a mask image Ilabel, usually
  %   created by bwlabel()
  % OUTPUT:
  % Each row in the output array OData represents
15 individual object
  % columns contain the following measurements:
  %
  %   1 - Index ("number" of an object);      8 -
  Solidity;
20 %   2 - X coordinate of the center of mass; 9 - Extent;
  %   3 - Y coordinate      "-"; 10 - Total
  Intensity;
  %   4 - Total Area (in pixels);      11 - Avg.
  Intensity;
25 %   5 - Ratio of MajorAxis/MinorAxis;      12 - Median
  Intensity;
  %   6 - Eccentricity;      13 - Intensity of
  20% bright pixel
  %   7 - EquivDiameter;      14 - Intensity of
30 80% bright pixel
  %
  % For details on morphological parameters see information
  on MatLab imfeature();

```

```
% Intensity parameters are either obvious or are
documented in comments in this file.
% Procedures in this file are documented in notebook file
"MATLAB Measuring Nuclei (1) 1-29-98.doc"

5
if (nargin ~= 2)
    error ('function requires exactly 2 parameters');
end
if (nargout ~= 1)
10    error ('function has 1 output argument (array X by
    14)');
end

% finished checking arguments

15
% first collect morphological parameters in a structure
array:
ImStats = imfeature(Ilabel, 'Area', 'Centroid',
    'MajorAxisLength',...
20    'MinorAxisLength', 'Eccentricity', 'EquivDiameter',
    ...
    'Solidity', 'Extent', 8 );

% now convert it into array (matrix) while collecting
25 intensity data for each object:

%preallocate output array:
numobjects = size(ImStats, 1);
OData = zeros(numobjects, 14);
30 %now convert ImStats into array and add intensity data to
    it
    for k=1:numobjects
        OData(k, 1) = k;
        OData(k, 2) = ImStats(k).Centroid(1);
```

```

        OData(k, 3) = ImStats(k).Centroid(2);
        OData(k, 4) = ImStats(k).Area;
        OData(k, 5) = (ImStats(k).MajorAxisLength) /
        (ImStats(k).MinorAxisLength);
5         OData(k, 6) = ImStats(k).Eccentricity ;
        OData(k, 7) = ImStats(k).EquivDiameter;
        OData(k, 8) = ImStats(k).Solidity;
        OData(k, 9) = ImStats(k).Extent;

10         % now collect and assign intensity parameters from
        image I

        object_pixels = find( Ilabel == k);
        object_area = size(object_pixels, 1); %same as total
15 number of pixels in the object
        object_intensities = double(I(object_pixels)); %
        need to convert to double to do math
        sorted_intensities = sort(object_intensities); %
        will need to get median, 20% and 80% pixels
20         total_intensity = sum(object_intensities, 1);
        avg_intensity = total_intensity / object_area;
        median_intensity = sorted_intensities( floor(
        object_area/2 ) + 1 );
        pix20 = sorted_intensities( floor(object_area*0.2)+1
25 ) ; %brightest pixel among dimmest 20%
        pix80 = sorted_intensities( floor(object_area*0.8)+1
        ) ;

        OData(k, 10) = total_intensity;
30         OData(k, 11) = avg_intensity;
        OData(k, 12) = median_intensity;
        OData(k, 13) = pix20; %brightest pixel among dimmest
        20%

```

```

        OData(k, 14) = pix80; %dimmet pixel among brightest
    20%
end %for

5  %===== end function
   GetObjectsData()=====

function Imask = MaskDNA1(I);
10 % MaskDNA1 - generates binary mask for cell nuclei
   through edge detection
   % EV 1-22-99; 2-6-99; 2-10-99
   % Imask = MaskDNA1(I)
   % PARAMETERS
15 %   I - intensity image (grayscale)
   % OUTPUT
   %   Imask - BW image with objects from I
   %
   % For more details see Notebook Matlab_DNA_masking1_1-22-
20 99.doc
   % Uses SDC Morphology Toolbox V0.7

   if (nargin ~= 1)
       error('Wrong number of input parameters');
25 end
   if (nargout ~= 1)
       error('Wrong number of output parameters: one output
       argument should be provided');
   end
30

   Imask = edge(I, 'canny');
   Imask = mmdil(Imask, mmsecross(1));
   Imask = mmero ( mmclohole(Imask,mmsecross(1)));

```

```

Imask = mmedgeoff(Imask, mmsecross(1));
% note that mmedgeoff this command removed FILLED OBJECTS
but not touching OUTLINES.
% these outlines can be removed by filtering:
5  Imask = medfilt2(Imask, [5 5]);

%=====end MaskDNA1 =====

```

Given the list of image files or montages of images as an input, this
 10 program creates an individual file for each image that contains the following
 quantitative measurements for all objects identified in the image:

- | | |
|---|------------------------------------|
| 1 - Index ("number" of an object); | 8 - Solidity; |
| 2 - X coordinate of the center of mass; | 9 - Extent; |
| 15 3 - Y coordinate "-"; | 10 - Total Intensity; |
| 4 - Total Area (in pixels); | 11 - Avg. Intensity; |
| 5 - Ratio of MajorAxis/MinorAxis; | 12 - Median Intensity; |
| 6 - Eccentricity; | 13 - Intensity of 20% bright pixel |
| 7 - EquivDiameter; | 14 - Intensity of 80% bright pixel |

20 A fragment of an output for a single file, containing 9 images of cells
 stained for DNA and acquired with a 10x objective. A montage image that was used
 as a source to generate data in A is presented in Fig. 17.

The same program also summarizes measurements across many files
 and performs statistical analysis of the summary data. It creates a summary file with
 25 the following data:

- | | |
|--|---------------------------------|
| 1 - Image file number; | |
| 2 - Average object Area (in pixels); | 3 - STD (standard deviation) of |
| 2; | |
| 30 4 - Avg. of Ratio of MajorAxis/MinorAxis; | 5 - STD of 4; |
| 6 - Avg. Eccentricity; | 7 - STD of 6; |
| 8 - Avg. EquivDiameter; | 9 - STD of 8; |
| 10 - Avg. of Solidity; | 11 - STD of 10; |

- | | | |
|---|---|----------------|
| | 12 - Avg. of Extent; | 13 - STD of 11 |
| | 14 - Avg. of objects Total Intensity; | 15 - STD of 14 |
| | 16 - Avg. of objects Avg Intensity; | 16 - STD of 15 |
| | 18 - Avg. of objects Median intensity; | 19 - STD of 18 |
| 5 | 20 - Avg. of objects intensity of 20% bright pixel; | 21 - STD of 19 |
| | 22 - Avg. of objects intensity of 80% bright pixel; | 23 - STD of 21 |

An example of summary output obtained by running AnalyseDNA against 10 montage files also is shown in Appendix B.

10 A seventh experiment was conducted in order to use sequence analysis algorithms to analyze features of cell images. In this experiment, HeLa cells were treated for 24 hours with several different compounds, and then fixed, and stained with a fluorescent DNA dye. One image of these cells was acquired for each of the treatments and morphometric parameters and features were measured:

15 Resulting measurements were arranged into a string of numbers and reduced to a pseudo- nucleic acid sequence using following rules: At any given position in the sequence a number was substituted by "t" (a code for thymidine) if its value is among highest 25% of the values at the corresponding position in the data set, "g" if it is between 50% and 25%, "c" if it is between 75% and 50%, and "a" if it
20 belongs to lowest 25% of values. Thus one descriptor or sequence was generated per treatment as illustrated in Fig. 18.

Resulting sequences were clustered using an AlignX module commercial software package Vector NTI (<http://informaxinc.com>), which uses a Neighbor Joining algorithm for sequence clustering.

25 The resulting dendrogram is presented in Fig 18. On the dendrogram the closest "leaves" correspond to the closest pseudo-sequences. Interestingly, compounds with similar mechanisms of action cluster together on the dendrogram. Another example of the generation of pseudo-sequences and clustering is shown in Fig. 19.

30 In some embodiments, techniques according to the present invention can provide tools for the later stages of drug development such as clinical trial design and patient management. The properties of known drugs such as clinical trial and patient response information will be used in a similar fashion as the pre-clinical

information to provide predictions about the properties of novel compounds. Because the human cell is the locus of drug action, a database containing drug-cell interactions can be able to provide predictive information for this aspect of drug development.

Although the above has generally described the present invention

5 according to specific systems, the present invention has a much broader range of applicability. In particular, the present invention is not limited to a particular kind of data about a cell, but can be applied to virtually any cellular data where an understanding about the workings of the cell is desired. Thus, in some embodiments, the techniques of the present invention could provide information about many

10 different types or groups of cells, substances, and genetic processes of all kinds. Of course, one of ordinary skill in the art would recognize other variations, modifications, and alternatives.

APPENDIX A

EV Table 1.doc

Example of the output of AnalysedDNA.m program
(measurements for a single 3 by 3 montage image)

File	Subimage	object#	X coord	Y coord	Area	Area ratio	Eccentricity	Equidiam	Solidity	Extant	Intensity	Avg. Intensity	Median	20% Intensity	80% Intensity
1	1	1	12.2697	132.453	165	1.17293	0.372654	13.3453	0.83567	0.736796	4405	31.7466	34	23	37
1	1	2	16.332	116.022	125	1.40394	0.764731	12.4137	0.965797	0.78123	4106	34.648	38	30	45
1	1	3	20.1073	75.2015	177	1.06845	0.437265	15.0121	0.817056	0.661406	4769	26.9425	28	22	31
1	1	4	21.6166	102.114	161	1.34515	0.471004	13.9826	0.814334	0.787857	3650	85.614	67	67	105
1	1	5	21.6166	102.114	161	1.34515	0.471004	13.9826	0.814334	0.787857	3650	85.614	67	67	105
1	1	6	22.622	259.334	204	2.33106	0.603209	14.1953	0.848889	0.713229	4502	66.8938	46	36	54
1	1	7	22.622	259.334	204	2.33106	0.603209	14.1953	0.848889	0.713229	4502	66.8938	46	36	54
1	1	8	23.629	167.572	166	1.34564	0.471456	10.4551	0.927083	0.711667	4223	47.4119	50	38	54
1	1	9	23.629	167.572	166	1.34564	0.471456	10.4551	0.927083	0.711667	4223	47.4119	50	38	54
1	1	10	27.144	366.021	41	1.84062	0.831512	7.73576	0.87037	0.632776	6667	141.1851	142	113	171
1	1	11	27.144	366.021	41	1.84062	0.831512	7.73576	0.87037	0.632776	6667	141.1851	142	113	171
1	1	12	34.0749	126.538	232	1.85101	0.531327	17.187	0.632941	0.70503	5632	42.2792	35	33	51
1	1	13	34.0749	126.538	232	1.85101	0.531327	17.187	0.632941	0.70503	5632	42.2792	35	33	51
1	1	14	32.7755	44.9622	197	2.27223	0.653301	13.7666	0.807407	0.704711	5110	21.215	21	26	39
1	1	15	32.7755	44.9622	197	2.27223	0.653301	13.7666	0.807407	0.704711	5110	21.215	21	26	39
1	1	16	36.0229	282.272	206	1.82382	0.554944	16.1653	0.82167	0.704711	5110	21.215	21	26	39
1	1	17	36.0229	282.272	206	1.82382	0.554944	16.1653	0.82167	0.704711	5110	21.215	21	26	39
1	1	18	48.1714	333.181	313	1.72885	0.818089	11.7245	0.721254	0.704711	5110	21.215	21	26	39
1	1	19	48.1714	333.181	313	1.72885	0.818089	11.7245	0.721254	0.704711	5110	21.215	21	26	39
1	1	20	71.6649	443.13	185	1.77876	0.409045	16.1653	0.82167	0.704711	5110	21.215	21	26	39
1	1	21	71.6649	443.13	185	1.77876	0.409045	16.1653	0.82167	0.704711	5110	21.215	21	26	39
1	1	22	76.7377	206.27	206	1.70117	0.432216	20.0747	0.82167	0.704711	5110	21.215	21	26	39
1	1	23	76.7377	206.27	206	1.70117	0.432216	20.0747	0.82167	0.704711	5110	21.215	21	26	39
1	1	24	81.7865	281.534	175	1.43132	0.511622	12.5138	0.82167	0.704711	5110	21.215	21	26	39
1	1	25	81.7865	281.534	175	1.43132	0.511622	12.5138	0.82167	0.704711	5110	21.215	21	26	39
1	1	26	65.1409	176.231	163	1.43132	0.511622	12.5138	0.82167	0.704711	5110	21.215	21	26	39
1	1	27	65.1409	176.231	163	1.43132	0.511622	12.5138	0.82167	0.704711	5110	21.215	21	26	39
1	1	28	61.7604	317.765	260	1.87333	0.652119	19.1922	0.9	0.66023	4543	27.0219	29	23	41
1	1	29	61.7604	317.765	260	1.87333	0.652119	19.1922	0.9	0.66023	4543	27.0219	29	23	41
1	1	30	96.6462	248.602	116	1.2774	0.422216	12.2373	0.82167	0.704711	5110	21.215	21	26	39
1	1	31	96.6462	248.602	116	1.2774	0.422216	12.2373	0.82167	0.704711	5110	21.215	21	26	39
1	1	32	103.933	93.2279	122	1.46415	0.728421	12.2373	0.82167	0.704711	5110	21.215	21	26	39
1	1	33	103.933	93.2279	122	1.46415	0.728421	12.2373	0.82167	0.704711	5110	21.215	21	26	39
1	1	34	105.47	155.407	134	1.3208	0.728421	12.2373	0.82167	0.704711	5110	21.215	21	26	39
1	1	35	105.47	155.407	134	1.3208	0.728421	12.2373	0.82167	0.704711	5110	21.215	21	26	39
1	1	36	125.322	170.145	152	1.3208	0.728421	12.2373	0.82167	0.704711	5110	21.215	21	26	39
1	1	37	125.322	170.145	152	1.3208	0.728421	12.2373	0.82167	0.704711	5110	21.215	21	26	39
1	1	38	126.86	307.383	266	1.75489	0.432208	13.3166	0.927432	0.734375	4429	45.3357	49	37	53
1	1	39	126.86	307.383	266	1.75489	0.432208	13.3166	0.927432	0.734375	4429	45.3357	49	37	53
1	1	40	130.102	411.5	164	1.19374	0.440319	16.4023	0.784073	0.536462	5810	34.7378	38	26	43
1	1	41	130.102	411.5	164	1.19374	0.440319	16.4023	0.784073	0.536462	5810	34.7378	38	26	43
1	1	42	132.439	352.545	167	1.39705	0.463212	15.4304	0.935	0.739113	5227	27.9519	27	22	34
1	1	43	132.439	352.545	167	1.39705	0.463212	15.4304	0.935	0.739113	5227	27.9519	27	22	34
1	1	44	134.455	209.056	101	1.15921	0.450346	11.3101	0.87069	0.617436	7293	72.2029	74	58	75
1	1	45	134.455	209.056	101	1.15921	0.450346	11.3101	0.87069	0.617436	7293	72.2029	74	58	75
1	1	46	140.833	102.008	121	1.47787	0.734306	12.4122	0.916667	0.732313	4806	38.7574	41	32	48
1	1	47	140.833	102.008	121	1.47787	0.734306	12.4122	0.916667	0.732313	4806	38.7574	41	32	48
1	1	48	144.664	59.8166	272	1.75519	0.821825	16.4087	0.937064	0.715544	5000	55.9004	54	41	64
1	1	49	144.664	59.8166	272	1.75519	0.821825	16.4087	0.937064	0.715544	5000	55.9004	54	41	64
1	1	50	147.063	438.435	161	1.3071	0.44397	14.2175	0.82167	0.704711	5110	21.215	21	26	39
1	1	51	147.063	438.435	161	1.3071	0.44397	14.2175	0.82167	0.704711	5110	21.215	21	26	39
1	1	52	155.488	256.224	224	1.08006	0.471892	17.7848	0.927432	0.734375	4429	45.3357	49	37	53
1	1	53	155.488	256.224	224	1.08006	0.471892	17.7848	0.927432	0.734375	4429	45.3357	49	37	53
1	1	54	160.875	362.254	141	1.39125	0.471892	17.7848	0.927432	0.734375	4429	45.3357	49	37	53
1	1	55	160.875	362.254	141	1.39125	0.471892	17.7848	0.927432	0.734375	4429	45.3357	49	37	53
1	1	56	165.412	134.247	232	1.71321	0.711055	15.8777	0.937064	0.715544	5000	55.9004	54	41	64
1	1	57	165.412	134.247	232	1.71321	0.711055	15.8777	0.937064	0.715544	5000	55.9004	54	41	64
1	1	58	174.014	281.534	166	1.41186	0.720416	14.6125	0.927432	0.734375	4429	45.3357	49	37	53
1	1	59	174.014	281.534	166	1.41186	0.720416	14.6125	0.927432	0.734375	4429	45.3357	49	37	53
1	1	60	176.184	310.214	127	1.13784	0.477004	12.3742	0.900763	0.755952	4671	39.5847	42	32	47
1	1	61	176.184	310.214	127	1.13784	0.477004	12.3742	0.900763	0.755952	4671	39.5847	42	32	47
1	1	62	182.4	392.474	147	1.34123	0.481604	12.7142	0.927432	0.734375	4429	45.3357	49	37	53
1	1	63	182.4	392.474	147	1.34123	0.481604	12.7142	0.927432	0.734375	4429	45.3357	49	37	53
1	1	64	186.316	242.319	166	1.31766	0.711857	14.7123	0.937064	0.715544	5000	55.9004	54	41	64
1	1	65	186.316	242.319	166	1.31766	0.711857	14.7123	0.937064	0.715544	5000	55.9004	54	41	64
1	1	66	200.742	92.7418	212	1.36363	0.683426	16.4882	0.937064	0.715544	5000	55.9004	54	41	64
1	1	67	200.742	92.7418	212	1.36363	0.683426	16.4882	0.937064	0.715544	5000	55.9004	54	41	64
1	1	68	199.188	156.725	91	1.04663	0.255161	10.7441	0.90039	0.752066	4988	66.022	47	36	54
1	1	69	199.188	156.725	91	1.04663	0.255161	10.7441	0.90039	0.752066	4988	66.022	47	36	54
1	1	70	206.47	86.871	264	1.96053	0.460182	18.134	0.916667	0.732313	4806	38.7574	41	32	48
1	1	71	206.47	86.871	264	1.96053	0.460182	18.134	0.916667	0.732313	4806	38.7574	41	32	48
1	1	72	208.752	10.0435	220	1.45129	0.605176	17.1127	0.946502	0.604198	5766	42.4809	44	34	52
1	1	73	208.752	10.0435	220	1.45129	0.605176	17.1127	0.946502	0.604198	5766	42.4809	44	34	52
1	1	74	212.314	369.465	167	1.37679	0.471567	15.6776	0.927432	0.734375	4429	45.3357	49	37	53
1	1	75	212.314	369.465	167	1.37679	0.471567	15.6776	0.927432	0.734375	442				

[illegible]

EV	Table	/doc
----	-------	------

EV Table 1.doc

60	223.098	346.61	122	1.19425	0.546784	12.5112	0.254612	0.732143	4532	36.4527	38	29	44
61	223.901	373.099	121	1.67159	0.645292	14.7555	0.271216	0.718238	4533	37.7527	35	28	40
62	230.327	380.872	118	1.64435	0.732026	15.0515	0.280092	0.718238	4534	38.4816	28	21	34
63	231.936	423.425	285	1.56216	0.805464	17.3115	0.316533	0.698933	4535	41.4864	26	26	39
64	241.415	435.715	480	1.27432	0.721432	17.3115	0.316533	0.698933	4536	41.4864	26	26	39
65	248.445	477.491	125	1.27432	0.721432	17.3115	0.316533	0.698933	4537	41.4864	26	26	39
66	248.795	477.491	125	1.27432	0.721432	17.3115	0.316533	0.698933	4538	41.4864	26	26	39
67	248.795	477.491	125	1.27432	0.721432	17.3115	0.316533	0.698933	4539	41.4864	26	26	39
68	248.795	477.491	125	1.27432	0.721432	17.3115	0.316533	0.698933	4540	41.4864	26	26	39
69	248.795	477.491	125	1.27432	0.721432	17.3115	0.316533	0.698933	4541	41.4864	26	26	39
70	251.401	459.551	147	1.34215	0.679415	14.0012	0.316533	0.698933	4542	41.4864	26	26	39
71	252.141	48.2734	139	1.28222	0.625913	13.4609	0.316533	0.698933	4543	41.4864	26	26	39
72	252.908	659.384	160	1.49214	0.721432	10.2273	0.316533	0.698933	4544	41.4864	26	26	39
73	253.935	302.612	201	1.40216	0.815321	15.9975	0.316533	0.698933	4545	41.4864	26	26	39
74	253.753	302.612	170	1.42161	0.839321	14.7123	0.316533	0.698933	4546	41.4864	26	26	39
75	262.19	24.4535	157	1.43319	0.721432	14.1386	0.316533	0.698933	4547	41.4864	26	26	39
76	304.73	201.073	176	1.13701	0.545623	11.0543	0.316533	0.698933	4548	41.4864	26	26	39
77	305.441	484.582	155	2.0845	0.877415	15.3476	0.316533	0.698933	4549	41.4864	26	26	39
78	307.247	40.3177	158	1.62279	0.839321	12.7162	0.316533	0.698933	4550	41.4864	26	26	39
79	308.274	325.239	127	1.62279	0.839321	12.7162	0.316533	0.698933	4551	41.4864	26	26	39
80	310.67	311.938	176	1.62279	0.839321	12.7162	0.316533	0.698933	4552	41.4864	26	26	39
81	315.691	217.246	229	1.52517	0.735029	17.0733	0.316533	0.698933	4553	41.4864	26	26	39
82	314.676	302.312	170	1.48119	0.712939	14.7123	0.316533	0.698933	4554	41.4864	26	26	39
83	320.441	216.309	228	1.57165	0.697326	17.0733	0.316533	0.698933	4555	41.4864	26	26	39
84	320.624	423.687	228	2.38169	0.951116	18.3552	0.316533	0.698933	4556	41.4864	26	26	39
85	325.638	376.28	216	2.38169	0.951116	18.3552	0.316533	0.698933	4557	41.4864	26	26	39
86	323.623	62.464	182	2.38169	0.951116	18.3552	0.316533	0.698933	4558	41.4864	26	26	39
87	320.016	216.309	228	2.38169	0.951116	18.3552	0.316533	0.698933	4559	41.4864	26	26	39
88	316.4	423.687	216	2.38169	0.951116	18.3552	0.316533	0.698933	4560	41.4864	26	26	39
89	311.04	423.687	216	2.38169	0.951116	18.3552	0.316533	0.698933	4561	41.4864	26	26	39
90	343.349	316.017	119	1.10502	0.423197	12.0619	0.316533	0.698933	4562	41.4864	26	26	39
91	351.021	351.021	219	1.49133	0.640967	11.0035	0.316533	0.698933	4563	41.4864	26	26	39
92	351.021	351.021	219	1.49133	0.640967	11.0035	0.316533	0.698933	4564	41.4864	26	26	39
93	351.021	351.021	219	1.49133	0.640967	11.0035	0.316533	0.698933	4565	41.4864	26	26	39
94	351.021	351.021	219	1.49133	0.640967	11.0035	0.316533	0.698933	4566	41.4864	26	26	39
95	355.603	326.309	110	1.53831	0.713668	11.3651	0.316533	0.698933	4567	41.4864	26	26	39
96	361.412	201.748	194	1.44194	0.720465	11.0343	0.316533	0.698933	4568	41.4864	26	26	39
97	361.412	201.748	194	1.44194	0.720465	11.0343	0.316533	0.698933	4569	41.4864	26	26	39
98	370.626	159.489	131	1.56713	0.765943	12.5112	0.316533	0.698933	4570	41.4864	26	26	39
99	373.976	497.422	223	1.64477	0.802332	16.9237	0.316533	0.698933	4571	41.4864	26	26	39
100	375.192	62.868	125	1.33274	0.673444	12.4157	0.316533	0.698933	4572	41.4864	26	26	39
101	375.192	62.868	125	1.33274	0.673444	12.4157	0.316533	0.698933	4573	41.4864	26	26	39
102	375.192	62.868	125	1.33274	0.673444	12.4157	0.316533	0.698933	4574	41.4864	26	26	39
103	375.192	62.868	125	1.33274	0.673444	12.4157	0.316533	0.698933	4575	41.4864	26	26	39
104	380.145	235.421	145	1.60064	0.737919	12.4157	0.316533	0.698933	4576	41.4864	26	26	39
105	381.421	207.214	184	1.73174	0.813914	14.0373	0.316533	0.698933	4577	41.4864	26	26	39
106	381.421	207.214	184	1.73174	0.813914	14.0373	0.316533	0.698933	4578	41.4864	26	26	39
107	401.246	248.112	119	1.50376	0.747781	10.2179	0.316533	0.698933	4579	41.4864	26	26	39
108	405.131	167.402	142	1.91382	0.855918	15.0947	0.316533	0.698933	4580	41.4864	26	26	39
109	405.131	167.402	142	1.91382	0.855918	15.0947	0.316533	0.698933	4581	41.4864	26	26	39
110	405.131	167.402	142	1.91382	0.855918	15.0947	0.316533	0.698933	4582	41.4864	26	26	39
111	405.131	167.402	142	1.91382	0.855918	15.0947	0.316533	0.698933	4583	41.4864	26	26	39
112	416.274	100.644	180	1.21602	0.587746	12.4655	0.316533	0.698933	4584	41.4864	26	26	39
113	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4585	41.4864	26	26	39
114	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4586	41.4864	26	26	39
115	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4587	41.4864	26	26	39
116	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4588	41.4864	26	26	39
117	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4589	41.4864	26	26	39
118	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4590	41.4864	26	26	39
119	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4591	41.4864	26	26	39
120	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4592	41.4864	26	26	39
121	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4593	41.4864	26	26	39
122	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4594	41.4864	26	26	39
123	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4595	41.4864	26	26	39
124	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4596	41.4864	26	26	39
125	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4597	41.4864	26	26	39
126	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4598	41.4864	26	26	39
127	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4599	41.4864	26	26	39
128	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4600	41.4864	26	26	39
129	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4601	41.4864	26	26	39
130	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4602	41.4864	26	26	39
131	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4603	41.4864	26	26	39
132	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4604	41.4864	26	26	39
133	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4605	41.4864	26	26	39
134	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4606	41.4864	26	26	39
135	426.611	225.589	170	1.85539	0.844701	17.3115	0.316533	0.698933	4607	41.4864	26	26	39

Page 4 of 16

EV Table 1.doc

1	136	445.46	266.449	289	2.80072	0.921085	19.5113	0.85521	0.68891	11063	41.0401	50	38	57
2	137	489.732	104.247	120	1.75179	0.821030	15.5536	0.855856	0.66667	8796	46.2421	50	39	57
3	138	517.287	117.287	125	1.65278	0.82085	15.575	0.813014	0.71381	5108	21.7333	29	22	31
4	139	556.076	145	131	1.51521	0.82085	15.575	0.813014	0.71381	5108	54	38	41	48
5	140	598.098	176.345	142	1.39192	0.82085	15.575	0.813014	0.71381	5108	36.0871	37	25	34
6	141	643.115	219.345	156	1.28174	0.82085	15.575	0.813014	0.71381	5108	35.0544	40	26	35
7	142	690.115	274.345	174	1.18974	0.82085	15.575	0.813014	0.71381	5108	33.2024	35	24	33
8	143	738.115	339.345	196	1.10974	0.82085	15.575	0.813014	0.71381	5108	31.2024	34	23	32
9	144	787.115	414.345	221	1.04974	0.82085	15.575	0.813014	0.71381	5108	29.2024	33	22	31
10	145	837.115	499.345	249	1.00974	0.82085	15.575	0.813014	0.71381	5108	27.2024	32	21	30
11	146	887.115	594.345	279	0.97974	0.82085	15.575	0.813014	0.71381	5108	25.2024	31	20	29
12	147	937.115	699.345	311	0.94974	0.82085	15.575	0.813014	0.71381	5108	23.2024	30	19	28
13	148	987.115	814.345	345	0.91974	0.82085	15.575	0.813014	0.71381	5108	21.2024	29	18	27
14	149	1037.115	939.345	381	0.88974	0.82085	15.575	0.813014	0.71381	5108	19.2024	28	17	26
15	150	1087.115	1074.345	419	0.85974	0.82085	15.575	0.813014	0.71381	5108	17.2024	27	16	25
16	151	1137.115	1219.345	459	0.82974	0.82085	15.575	0.813014	0.71381	5108	15.2024	26	15	24
17	152	1187.115	1374.345	501	0.79974	0.82085	15.575	0.813014	0.71381	5108	13.2024	25	14	23
18	153	1237.115	1539.345	545	0.76974	0.82085	15.575	0.813014	0.71381	5108	11.2024	24	13	22
19	154	1287.115	1714.345	591	0.73974	0.82085	15.575	0.813014	0.71381	5108	9.2024	23	12	21
20	155	1337.115	1899.345	639	0.70974	0.82085	15.575	0.813014	0.71381	5108	7.2024	22	11	20
21	156	1387.115	2094.345	689	0.67974	0.82085	15.575	0.813014	0.71381	5108	5.2024	21	10	19
22	157	1437.115	2299.345	741	0.64974	0.82085	15.575	0.813014	0.71381	5108	3.2024	20	9	18
23	158	1487.115	2514.345	795	0.61974	0.82085	15.575	0.813014	0.71381	5108	1.2024	19	8	17
24	159	1537.115	2739.345	851	0.58974	0.82085	15.575	0.813014	0.71381	5108	0.2024	18	7	16
25	160	1587.115	2974.345	909	0.55974	0.82085	15.575	0.813014	0.71381	5108	0.2024	17	6	15
26	161	1637.115	3219.345	969	0.52974	0.82085	15.575	0.813014	0.71381	5108	0.2024	16	5	14
27	162	1687.115	3474.345	1031	0.49974	0.82085	15.575	0.813014	0.71381	5108	0.2024	15	4	13
28	163	1737.115	3739.345	1095	0.46974	0.82085	15.575	0.813014	0.71381	5108	0.2024	14	3	12
29	164	1787.115	4014.345	1161	0.43974	0.82085	15.575	0.813014	0.71381	5108	0.2024	13	2	11
30	165	1837.115	4299.345	1229	0.40974	0.82085	15.575	0.813014	0.71381	5108	0.2024	12	1	10
31	166	1887.115	4594.345	1299	0.37974	0.82085	15.575	0.813014	0.71381	5108	0.2024	11	0	9
32	167	1937.115	4899.345	1371	0.34974	0.82085	15.575	0.813014	0.71381	5108	0.2024	10	0	8
33	168	1987.115	5214.345	1445	0.31974	0.82085	15.575	0.813014	0.71381	5108	0.2024	9	0	7
34	169	2037.115	5539.345	1521	0.28974	0.82085	15.575	0.813014	0.71381	5108	0.2024	8	0	6
35	170	2087.115	5874.345	1599	0.25974	0.82085	15.575	0.813014	0.71381	5108	0.2024	7	0	5
36	171	2137.115	6219.345	1679	0.22974	0.82085	15.575	0.813014	0.71381	5108	0.2024	6	0	4
37	172	2187.115	6574.345	1761	0.19974	0.82085	15.575	0.813014	0.71381	5108	0.2024	5	0	3
38	173	2237.115	6939.345	1845	0.16974	0.82085	15.575	0.813014	0.71381	5108	0.2024	4	0	2
39	174	2287.115	7314.345	1931	0.13974	0.82085	15.575	0.813014	0.71381	5108	0.2024	3	0	1
40	175	2337.115	7699.345	2019	0.10974	0.82085	15.575	0.813014	0.71381	5108	0.2024	2	0	0
41	176	2387.115	8094.345	2109	0.07974	0.82085	15.575	0.813014	0.71381	5108	0.2024	1	0	0
42	177	2437.115	8499.345	2201	0.04974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
43	178	2487.115	8914.345	2295	0.01974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
44	179	2537.115	9339.345	2391	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
45	180	2587.115	9774.345	2489	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
46	181	2637.115	10219.345	2589	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
47	182	2687.115	10674.345	2691	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
48	183	2737.115	11139.345	2795	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
49	184	2787.115	11614.345	2901	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
50	185	2837.115	12099.345	3009	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
51	186	2887.115	12594.345	3119	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
52	187	2937.115	13099.345	3231	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
53	188	2987.115	13614.345	3345	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
54	189	3037.115	14139.345	3461	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
55	190	3087.115	14674.345	3579	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
56	191	3137.115	15219.345	3699	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
57	192	3187.115	15774.345	3821	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
58	193	3237.115	16339.345	3945	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
59	194	3287.115	16914.345	4071	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
60	195	3337.115	17499.345	4209	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
61	196	3387.115	18094.345	4349	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
62	197	3437.115	18699.345	4491	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
63	198	3487.115	19314.345	4635	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
64	199	3537.115	19939.345	4781	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
65	200	3587.115	20574.345	4929	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
66	201	3637.115	21219.345	5079	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
67	202	3687.115	21874.345	5231	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
68	203	3737.115	22539.345	5385	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
69	204	3787.115	23214.345	5541	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
70	205	3837.115	23899.345	5699	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
71	206	3887.115	24594.345	5859	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
72	207	3937.115	25299.345	6021	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
73	208	3987.115	26014.345	6185	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
74	209	4037.115	26739.345	6351	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
75	210	4087.115	27474.345	6519	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
76	211	4137.115	28219.345	6689	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
77	212	4187.115	28974.345	6861	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
78	213	4237.115	29739.345	7035	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
79	214	4287.115	30514.345	7211	0.00974	0.82085	15.575	0.813014	0.71381	5108	0.2024	0	0	0
80	215	4337.115	31299.345	7389	0.00974	0.82085	15.5							

EV Table 1.doc

1	3	74	270.921	432.427	177	1.37886	0.424112	19.0121	0.317088	0.743697	9749	55.0791	55	42	69
2	3	75	232.484	437.243	221	1.32188	0.615964	16.7146	0.484	0.613869	9572	55.0791	55	42	69
3	3	76	232.822	345.879	208	2.53252	0.91876	16.2327	0.508297	0.519048	9910	51.0102	32	33	57
4	3	77	267.91	395.164	488	2.0161	0.872636	24.3267	0.711641	0.530435	15133	51.0102	32	24	28
5	3	78	282.431	390.639	164	1.86664	0.841395	13.4006	0.9	0.466667	7827	54.3842	36	42	48
6	3	79	288.331	390.639	216	2.86567	0.928094	13.4006	0.716125	0.322109	10486	54.3842	36	30	32
7	3	80	280.063	492.403	189	1.73762	0.989179	15.5126	0.211551	0.75	772	25.0322	47	20	39
8	3	81	282.268	27.4071	112	1.49137	0.741848	15.5126	0.488889	0.427222	1842	45.5464	47	36	34
9	3	82	286.84	279.133	232	1.2274	0.521128	15.5126	0.888889	0.427222	1842	45.5464	47	36	34
10	3	83	286.587	44.0877	114	1.25651	0.717139	15.5126	0.989179	0.490909	4589	40.2314	40	31	49
11	3	84	294.161	201.421	124	1.30511	0.651525	15.5126	0.739134	0.428642	6013	40.2314	40	31	49
12	3	85	294.161	201.421	124	1.30511	0.651525	15.5126	0.907089	0.716667	6258	40.2314	40	31	49
13	3	86	299.08	372.084	175	1.32743	0.681635	17.2409	0.42871	0.724456	5552	36.347	38	28	44
14	3	87	321.781	347.283	236	1.40716	0.781031	17.2409	0.494625	0.428079	5032	44.708	47	36	51
15	3	88	308.432	376.07	128	1.37652	0.487138	17.2409	0.934858	0.628079	10710	43	47	35	51
16	3	89	317.311	458.408	473	1.32602	0.581645	12.7442	0.925536	0.740123	4466	36.4094	38	28	44
17	3	90	310.433	41.0155	223	1.44555	0.794166	10.2795	0.927805	0.421154	5695	30.0158	29	21	43
18	3	91	319.03	358.212	99	1.48285	0.742184	24.5006	0.632182	0.615865	16173	34.1846	35	23	42
19	3	92	310.33	358.212	100	1.48285	0.742184	24.5006	0.909301	0.712786	4873	46.75	51	38	58
20	3	93	320.597	216.664	134	1.40176	0.700768	13.0619	0.909301	0.407407	4791	43.9541	46	34	54
21	3	94	319.03	358.212	99	1.48285	0.742184	24.5006	0.909301	0.407407	4791	43.9541	46	34	54
22	3	95	320.597	216.664	134	1.40176	0.700768	13.0619	0.909301	0.407407	4791	43.9541	46	34	54
23	3	96	320.597	216.664	134	1.40176	0.700768	13.0619	0.909301	0.407407	4791	43.9541	46	34	54
24	3	97	337.319	317.873	271	2.09271	0.878442	19.3755	0.925536	0.413121	4913	36.4791	37	28	44
25	3	98	337.319	317.873	271	2.09271	0.878442	19.3755	0.925536	0.413121	4913	36.4791	37	28	44
26	3	99	331.564	304.017	172	1.40963	0.783119	15.4552	0.909301	0.413121	4913	36.4791	37	28	44
27	3	100	338.308	300.432	133	1.45882	0.501154	15.4552	0.909301	0.413121	4913	36.4791	37	28	44
28	3	101	342.452	300.068	118	1.40229	0.701558	12.3841	0.909301	0.413121	4913	36.4791	37	28	44
29	3	102	349.47	287.802	161	1.78123	0.828816	12.3841	0.909301	0.413121	4913	36.4791	37	28	44
30	3	103	355.573	324.459	267	1.21295	0.81317	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
31	3	104	342.162	21.4599	127	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
32	3	105	342.162	21.4599	127	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
33	3	106	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
34	3	107	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
35	3	108	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
36	3	109	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
37	3	110	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
38	3	111	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
39	3	112	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
40	3	113	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
41	3	114	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
42	3	115	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
43	3	116	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
44	3	117	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
45	3	118	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
46	3	119	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
47	3	120	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
48	3	121	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
49	3	122	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
50	3	123	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
51	3	124	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
52	3	125	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
53	3	126	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
54	3	127	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
55	3	128	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
56	3	129	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
57	3	130	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
58	3	131	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
59	3	132	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
60	3	133	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
61	3	134	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
62	3	135	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
63	3	136	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
64	3	137	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
65	3	138	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
66	3	139	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
67	3	140	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
68	3	141	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
69	3	142	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
70	3	143	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
71	3	144	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
72	3	145	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
73	3	146	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
74	3	147	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
75	3	148	344.781	38.3194	114	1.51295	0.717139	14.3755	0.909301	0.413121	4913	36.4791	37	28	44
76	3	149	344.781	38.3194	114	1.51295	0.717139								

EV Table 1.doc

1	1	150	481.886	386.084	24	1.18938	0.341385	6.47538	0.945946	0.033333	3545	101.296	105	88	117
1	1	151	487.258	374.769	120	2.04206	0.871881	12.8653	0.902778	0.037255	4853	35.0231	16	26	46
1	1	152	491.926	364.508	128	1.24055	0.591944	12.7662	0.920463	0.041111	5156	37.2975	26	30	46
1	1	153	501.436	355.432	39	1.4308	0.715013	7.04773	0.905977	0.04125	3460	31.7856	35	39	59
1	1	154	512.431	347.431	152	1.3561	0.675468	13.5116	0.923212	0.040745	4322	41.1726	50	57	77
1	1	155	524.201	340.043	113	1.35777	0.692337	12.8756	0.923215	0.040745	4311	39.3261	40	48	68
1	1	156	537.151	333.043	115	1.35777	0.692337	12.8756	0.923215	0.040745	4311	39.3261	40	48	68
1	1	157	551.376	327.042	137	1.66619	0.800432	12.2075	0.923215	0.040745	4311	39.3261	40	48	68
1	1	158	566.402	321.245	151	1.66619	0.800432	12.2075	0.923215	0.040745	4311	39.3261	40	48	68
1	1	159	582.029	315.559	102	1.26259	0.624651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	160	598.376	310.043	202	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	161	615.376	304.77	192	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	162	632.976	299.504	119	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	163	651.076	294.278	259	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	164	669.776	289.043	184	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	165	689.076	283.77	166	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	166	708.976	278.504	172	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	167	729.476	273.278	81	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	168	750.576	268.043	102	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	169	772.076	262.77	100	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	170	794.076	257.504	159	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	171	816.576	252.278	259	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	172	839.576	247.043	184	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	173	863.076	241.77	166	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	174	887.076	236.504	172	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	175	911.576	231.278	81	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	176	936.576	226.043	102	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	177	962.076	220.77	100	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	178	988.076	215.504	159	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	179	1014.576	210.278	259	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	180	1041.576	205.043	184	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	181	1069.076	199.77	166	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	182	1097.076	194.504	172	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	183	1125.576	189.278	81	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	184	1154.576	184.043	102	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	185	1184.076	178.77	100	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	186	1214.076	173.504	159	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	187	1244.576	168.278	259	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	188	1275.576	163.043	184	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	189	1307.076	157.77	166	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	190	1339.076	152.504	172	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	191	1371.576	147.278	81	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	192	1404.576	142.043	102	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	193	1438.076	136.77	100	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	194	1472.076	131.504	159	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	195	1506.576	126.278	259	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	196	1541.576	121.043	184	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	197	1577.076	115.77	166	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	198	1613.076	110.504	172	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	199	1649.576	105.278	81	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	200	1686.576	100.043	102	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	201	1724.076	94.77	100	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	202	1762.076	89.504	159	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	203	1800.576	84.278	259	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	204	1839.576	79.043	184	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	205	1879.076	73.77	166	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	206	1919.076	68.504	172	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	207	1959.576	63.278	81	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	208	2000.576	58.043	102	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	209	2042.076	52.77	100	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	210	2084.076	47.504	159	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	211	2126.576	42.278	259	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	212	2169.576	37.043	184	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	213	2213.076	31.77	166	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	214	2257.076	26.504	172	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	215	2301.576	21.278	81	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	216	2346.576	16.043	102	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	217	2392.076	10.77	100	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	218	2438.076	5.504	159	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	219	2484.576	0.278	259	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	220	2531.576	-4.999	184	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	221	2579.076	-10.273	166	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	222	2626.576	-15.547	172	2.26259	0.824651	11.8497	0.905977	0.040745	4311	39.3261	40	48	68
1	1	223	2674.076	-20.821	81	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	224	2721.576	-26.095	102	1.4308	0.715013	7.04773	0.905977	0.040745	4311	39.3261	40	48	68
1	1	225	2769.076	-31.369	100	1.4308	0.715013	7.04773	0.9059						

Page 8 of 16

[illegible]

[illegible]

EV Table 1.doc

1	104	335.018	312.36	114	1.70064	0.923112	12.0478	0.897639	0.653734	4181	34.4734	38	29	41
2	105	361.744	169.021	129	1.40316	0.781040	12.9155	0.908451	0.661338	1536	58.4196	40	46	71
3	106	370.42	168.766	132	1.37346	0.64247	11.9416	0.903226	0.717808	4218	37.4607	40	30	51
4	107	379.464	161.258	137	1.31571	0.548772	11.1122	0.895148	0.731809	8130	42.7216	41	36	55
5	108	382.727	164.484	141	1.24828	0.420259	15.4304	0.912195	0.611111	8130	44.1497	49	38	61
6	109	385.96	157.146	147	1.20204	0.333333	19.6116	0.907188	0.611111	11210	36.9987	39	39	81
7	110	389.152	157.146	151	1.16233	0.221985	14.8443	0.915189	0.735	8757	50.3216	32	27	61
8	111	392.336	157.146	156	1.12554	0.121333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
9	112	395.52	157.146	161	1.08881	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
10	113	398.704	157.146	166	1.05208	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
11	114	401.886	157.146	171	1.01535	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
12	115	405.068	157.146	176	0.97862	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
13	116	408.250	157.146	181	0.94189	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
14	117	411.432	157.146	186	0.90516	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
15	118	414.614	157.146	191	0.86843	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
16	119	417.796	157.146	196	0.83170	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
17	120	420.978	157.146	201	0.79497	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
18	121	424.160	157.146	206	0.75824	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
19	122	427.342	157.146	211	0.72151	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
20	123	430.524	157.146	216	0.68478	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
21	124	433.706	157.146	221	0.64805	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
22	125	436.888	157.146	226	0.61132	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
23	126	440.070	157.146	231	0.57459	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
24	127	443.252	157.146	236	0.53786	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
25	128	446.434	157.146	241	0.50113	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
26	129	449.616	157.146	246	0.46440	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
27	130	452.798	157.146	251	0.42767	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
28	131	455.980	157.146	256	0.39094	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
29	132	459.162	157.146	261	0.35421	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
30	133	462.344	157.146	266	0.31748	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
31	134	465.526	157.146	271	0.28075	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
32	135	468.708	157.146	276	0.24402	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
33	136	471.890	157.146	281	0.20729	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
34	137	475.072	157.146	286	0.17056	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
35	138	478.254	157.146	291	0.13383	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
36	139	481.436	157.146	296	0.09710	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
37	140	484.618	157.146	301	0.06037	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
38	141	487.800	157.146	306	0.02364	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
39	142	490.982	157.146	311	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
40	143	494.164	157.146	316	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
41	144	497.346	157.146	321	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
42	145	500.528	157.146	326	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
43	146	503.710	157.146	331	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
44	147	506.892	157.146	336	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
45	148	510.074	157.146	341	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
46	149	513.256	157.146	346	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
47	150	516.438	157.146	351	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
48	151	519.620	157.146	356	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
49	152	522.802	157.146	361	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
50	153	525.984	157.146	366	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
51	154	529.166	157.146	371	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
52	155	532.348	157.146	376	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
53	156	535.530	157.146	381	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
54	157	538.712	157.146	386	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
55	158	541.894	157.146	391	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
56	159	545.076	157.146	396	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
57	160	548.258	157.146	401	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
58	161	551.440	157.146	406	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
59	162	554.622	157.146	411	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
60	163	557.804	157.146	416	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
61	164	560.986	157.146	421	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
62	165	564.168	157.146	426	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
63	166	567.350	157.146	431	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
64	167	570.532	157.146	436	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
65	168	573.714	157.146	441	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
66	169	576.896	157.146	446	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
67	170	580.078	157.146	451	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
68	171	583.260	157.146	456	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
69	172	586.442	157.146	461	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
70	173	589.624	157.146	466	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
71	174	592.806	157.146	471	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
72	175	595.988	157.146	476	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
73	176	599.170	157.146	481	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
74	177	602.352	157.146	486	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
75	178	605.534	157.146	491	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
76	179	608.716	157.146	496	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
77	180	611.898	157.146	501	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
78	181	615.080	157.146	506	0.00000	0.021333	13.158	0.897337	0.635646	4377	23.4514	35	32	61
79	182	618.262	157.146	511	0.00000	0.021333	13							

Page 12 of 16

1	1	11.1154	287.587	104	1.42586	0.712037	11.5073	0.504318	0.473259	3325	35.0788	29	31
1	2	9.0035	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	3	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	4	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	5	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	6	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	7	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	8	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	9	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	10	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	11	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	12	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	13	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	14	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	15	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	16	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	17	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	18	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	19	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	20	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	21	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	22	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	23	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	24	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	25	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	26	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	27	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	28	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	29	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	30	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	31	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	32	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	33	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	34	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	35	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	36	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	37	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	38	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	39	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	40	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	41	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	42	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	43	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	44	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	45	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	46	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	47	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	48	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	49	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	50	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	51	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	52	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	53	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	54	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	55	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	56	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	57	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	58	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	59	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	60	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	61	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	62	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	63	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	64	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	65	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	66	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	67	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	68	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	69	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	70	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	71	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	72	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	73	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	74	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	75	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43
1	76	11.1154	425.712	67	1.2659	0.538931	9.1817	0.48	0.732332	2055	45.2165	48	43

EV Table 1.doc

Page 13 of 16

Page 14 of 16

EV Table 1.doc

1	1	21.2188	346.125	64	1.20161	0.630312	9.07303	0.476712	0.727273	3642	51.0063	57	44
2	1	34.0233	432.916	129	1.24546	0.672231	12.0159	0.521429	0.661538	3872	30.0155	30	23
3	1	39.6228	516.229	140	1.25761	0.701463	13.3512	0.509091	0.701947	4313	30.0071	30	23
4	1	39.9272	531.316	208	1.24789	0.70403	14.1953	0.519443	0.677332	5011	43.7533	43	33
5	1	38.3362	528.343	178	1.05005	0.701765	15.0549	0.527083	0.701131	9196	51.6589	52	36
6	1	36.3667	513.346	120	1.17123	0.527049	10.8198	0.520245	0.761306	4709	31.3933	33	26
7	1	38.3745	542.721	135	1.17168	0.527265	10.4031	0.524737	0.761323	3860	46.4235	49	29
8	1	38.3745	542.721	135	1.40446	0.702172	15.3034	0.524621	0.682824	4279	30.7602	32	28
9	1	45.7156	703.948	189	1.77145	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
10	1	55.8006	62.5374	146	1.41539	0.785475	13.4243	0.521654	0.677857	4110	23.3869	24	22
11	1	55.8006	62.5374	146	1.31697	0.630312	12.666	0.5	0.625208	4098	52.3736	32	25
12	1	65.4198	126.373	126	1.41539	0.630312	20.3108	0.79601	0.625208	4098	52.3736	32	25
13	1	65.4198	126.373	126	1.455339	0.760018	15.757	0.519911	0.625142	18200	46.7621	48	35
14	1	61.6515	292.139	195	1.455339	0.760018	15.757	0.519911	0.625142	18200	46.7621	48	35
15	1	61.6515	292.139	195	1.77168	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
16	1	71.2222	315.178	45	1.42638	0.716145	17.5954	0.501823	0.716145	3166	74.9	74	54
17	1	71.2222	315.178	45	1.45506	0.730839	13.9572	0.519911	0.625142	18200	46.7621	48	35
18	1	71.2222	315.178	45	1.45506	0.730839	13.9572	0.519911	0.625142	18200	46.7621	48	35
19	1	76.1075	362.052	93	1.46164	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
20	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
21	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
22	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
23	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
24	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
25	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
26	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
27	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
28	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
29	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
30	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
31	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
32	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
33	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
34	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
35	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
36	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
37	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
38	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
39	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
40	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
41	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
42	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
43	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
44	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
45	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
46	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
47	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
48	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
49	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
50	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
51	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
52	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
53	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
54	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
55	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
56	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
57	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
58	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
59	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
60	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
61	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
62	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
63	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
64	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
65	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
66	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
67	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
68	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
69	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
70	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
71	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
72	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
73	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
74	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
75	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
76	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
77	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
78	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
79	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
80	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
81	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22
82	1	81.9478	91.3618	192	1.27679	0.625426	15.3034	0.521654	0.677857	4110	23.3869	24	22

EV Table 1.doc

1	283.416	101.811	127	1.4866	0.405787	12.7162	0.913669	0.461439	4425	38.4173	37	29	44
2	284.655	410.471	133	1.53854	0.755964	34.8445	0.913134	0.759772	7870	44.3333	41	35	51
3	285.45	289.912	64	1.31654	0.758214	10.2118	0.902226	0.460134	4311	50.131	31	39	62
4	286.312	60.1601	154	1.13337	0.476646	14.0028	0.816657	0.713333	4326	22.1886	30	23	36
5	287.889	305.311	90	1.08786	0.357114	10.7647	0.895951	0.713302	4136	45.9332	46	36	54
6	289.374	47.8166	107	1.27121	0.645038	11.472	0.89816	0.716286	4057	38.0093	39	39	44
7	290.3	332.155	106	1.53544	0.758638	11.4174	0.93053	0.818327	4300	44.2394	44	36	54
8	291.561	348.784	118	2.38788	0.908058	13.7733	0.840809	0.328371	8947	60.4527	40	41	54
9	293.159	364.741	44	1.28753	0.437205	7.40882	0.866154	0.6975	7440	85.4163	46	45	50
10	294.183	388.246	224	2.05865	0.874094	18.888	0.912886	0.39491	9246	41.2784	43	42	50
11	295.237	742.353	359	2.13597	0.848434	21.3797	0.810384	0.634611	13917	55.2008	36	42	64
12	296.322	322.118	182	1.46417	0.738234	15.2227	0.928571	0.716462	9774	32.0349	35	41	64
13	297.45	51.4724	163	1.82189	0.435602	14.4062	0.910615	0.632116	4364	26.775	32	39	52
14	298.5	356.403	158	1.26078	0.428164	14.1935	0.923977	0.732111	8946	46.1452	32	39	52
15	299.73	423.568	241	1.46335	0.739238	17.3172	0.848212	0.725463	4249	28.1391	29	32	34
16	300.921	425.407	152	1.33618	0.462277	14.7502	0.855732	0.646467	5659	202.107	21	166	238
17	302.143	325.407	152	1.33618	0.462277	14.7502	0.855732	0.646467	5659	202.107	21	166	238
18	303.425	386.534	145	1.40948	0.739153	14.5475	0.935135	0.738712	4599	23.1616	22	21	33
19	304.72	38.9614	217	1.37417	0.493884	13.5475	0.925487	0.70339	8423	58.0497	40	47	69
20	306.06	68.4019	74	2.38778	0.916421	16.4211	0.907193	0.645013	4555	21.1731	21	17	24
21	307.454	137.455	11	1.13876	0.470293	2.74241	0.844118	0.650308	7203	98.8028	99	76	114
22	308.864	213.176	148	1.33856	0.433143	13.7273	0.845571	0.6171	210	19.0909	19	16	22
23	310.312	102.314	166	2.28147	0.99901	14.5381	0.922232	0.65873	4596	27.4887	29	22	33
24	311.777	170.057	404	1.31694	0.450688	23.4601	0.920876	0.708294	20716	51.2732	52	38	44
25	313.190	243.499	328	1.4413	0.739718	20.4358	0.821818	0.788662	12022	38.6324	37	27	44
26	314.638	462.116	69	1.52746	0.71603	9.37302	0.896104	0.427273	3544	51.2624	32	41	62
27	316.101	691.639	135	1.6691	0.900653	14.0483	0.922419	0.65821	5726	28.9113	28	29	38
28	317.592	297.197	213	1.57725	0.431707	15.757	0.924171	0.716286	5007	28.7312	28	35	54
29	319.112	406.037	455.366	1.57313	0.733937	10.2179	0.811111	0.739238	2845	46.1897	31	38	58
30	320.672	217.136	124	1.23258	0.392299	7.40239	0.848983	0.716286	5007	46.1897	31	38	58
31	322.282	404.581	72	1.45318	0.728129	12.3451	0.913849	0.716286	4216	82.1094	51	53	101
32	323.938	243.171	216	1.14162	0.471379	9.57441	0.818048	0.716286	4216	32.3	35	36	46
33	325.643	288.014	220	2.13928	0.433154	20.2185	0.813602	0.716286	4216	32.3	35	36	46
34	327.398	609.116	230	2.13928	0.433154	20.2185	0.813602	0.716286	4216	32.3	35	36	46
35	329.153	57.6413	145	1.36234	0.820706	17.1137	0.922381	0.740792	9481	51.9464	32	41	61
36	330.907	437.769	92	1.40161	0.708756	14.1368	0.816532	0.742857	3333	34.1211	35	42	61
37	332.650	338.483	161	2.28722	0.823708	12.3092	0.901515	0.708123	4146	27.2733	27	20	33
38	334.393	624.193	119	1.28823	0.437708	12.3092	0.901515	0.708123	4146	27.2733	27	20	33
39	336.136	624.193	119	1.28823	0.437708	12.3092	0.901515	0.708123	4146	27.2733	27	20	33
40	337.879	449.5	44	1.13755	0.503805	13.7662	0.920863	0.737396	4293	32.9381	35	26	40
41	339.622	449.5	44	1.23615	0.58786	7.48422	0.88	0.737396	4293	32.9381	35	26	40
42	341.365	407.954	130	1.44995	0.709392	12.8453	0.913193	0.716286	7021	54.0077	34	41	66
43	343.108	187.428	132	1.44995	0.709392	12.8453	0.913193	0.716286	7021	54.0077	34	41	66
44	344.851	215.246	121	1.27349	0.61915	12.4123	0.902995	0.716286	4943	32.3724	33	28	43
45	346.594	10.1837	49	1.4709	0.733302	7.89845	0.907407	0.716286	5891	25.4924	31	28	43
46	348.337	132.365	160	1.4603	0.702675	14.6255	0.913232	0.717778	4890	120.124	122	92	137
47	350.080	230.344	160	1.45375	0.48109	14.713	0.816406	0.716286	4890	25.4924	31	28	43
48	351.823	41.2345	145	1.47223	0.717238	15.5945	0.913193	0.716286	4295	25.4924	31	28	43
49	353.566	311.423	171	2.14813	0.845034	16.4211	0.921184	0.702105	9350	40.9519	30	38	60
50	355.309	624.193	119	1.27169	0.437708	12.3092	0.901515	0.708123	4146	27.2733	27	20	33
51	357.052	338.483	161	2.13112	0.818453	15.0543	0.922238	0.708123	4146	27.2733	27	20	33

EV Table 2.doc

Example of the summary output of AnalyseDNA.m program
(summary for 10 3 by 3 montage images)

1	1197	165.812	79.3518	1.53849	0.726461	0.137996	16.0612	3.319	0.903373	0.0330365	0.701248	0.071176	649.26	3496.35	61.539	19.352	42.9466	19.9393
2	1205	14.334	50.9906	22.4584	0.399942	0.727935	0.138462	14.2634	3.43288	0.806571	0.0341453	0.70177	0.0730289	8784.37	3095.1	62.2416	17.0965	43.8245
3	1359	164.57	80.0679	1.9728	0.397935	0.72552	0.13163	14.0853	3.23966	0.803267	0.0341822	0.702891	0.0710005	6881.06	3525.81	64.7167	20.5918	44.3818
4	1368	172.015	59.7145	1.90189	0.389482	0.737142	0.142518	14.7351	3.41812	0.802766	0.0374254	0.693945	0.0733889	6997.68	6212.87	63.1798	19.8582	44.9561
5	1418	171.921	50.4617	1.91819	0.400193	0.721044	0.141704	14.738	3.45065	0.804152	0.0379801	0.70023	0.0734884	7030.22	6153.04	64.0559	21.4761	45.8522
6	1418	146.358	52.2517	1.91819	0.425512	0.728414	0.132723	14.0974	3.38686	0.804614	0.0352731	0.696204	0.0782855	6813.2	3924.12	64.261	19.3396	45.9405
7	1758	175.864	53.8055	22.8172	0.451022	0.694813	0.185467	11.724	5.27441	0.893111	0.081729	0.704526	0.0892393	5182.51	4392.13	74.9743	21.1949	36.16
8	1760	171.987	84.1244	1.93101	0.405167	0.723421	0.137899	14.3833	3.40593	0.806384	0.0357179	0.703574	0.0751602	6865.87	3787.48	62.2732	17.2223	43.9781
9	1770	166.53	84.5094	1.49349	0.404886	0.726461	0.139548	14.1402	3.47784	0.803208	0.0358298	0.700331	0.0766773	6576.54	4022.38	61.4064	17.8994	43.143
10	1805	159.406	82.618	1.97884	0.400372	0.717147	0.139005	13.8295	3.48936	0.804278	0.0378113	0.702759	0.07572	6587.18	3788.39	64.2141	20.7694	45.8765

CLAIMS

What is claimed is:

1. A method of predicting a property of a manipulation of cells based
5 upon a descriptor, said method comprising:
 providing a plurality of cells;
 manipulating said plurality of cells;
 capturing a morphological value from said plurality of cells;
 assigning a degree of presence of said morphological value; and
10 storing said morphological value and said degree of presence;
 wherein said descriptor is derived from a first component of a cell and
a second component of said cell, said capturing said morphometric value from said
plurality of cells comprises determining a relationship between said first component
and said second component.
- 15 2. The method of claim 1 wherein said first component and said second
component are selected from a protein, a protein modification, a nucleic acid, a lipid,
a carbohydrate, a subcellular structure and an organelle.
3. The method of 1 wherein said step of manipulation occurs in a manner
selected from a electrical source, a chemical source, a thermal source, a gravitational
20 source, a nuclear source, a temporal source, and a biological source
4. The method of claim 3 wherein said chemical source is selected from a
pharmacological candidate and a drug screening library.
5. The method of claim 1 wherein said morphological value is selected
from a count, an area, a perimeter, a length, a breadth, a fiber length, a fiber breadth, a
25 shape factor, a elliptical form factor, an inner radius, an outer radius, a mean radius,
an equivalent radius, an equivalent sphere volume, an equivalent prolate volume, an
equivalent oblate volume, an equivalent sphere surface area, an average gray value, a
total gray value, and an optical density.
6. The method of claim 1 wherein said degree of presence is
30 multiple of a quantized value.

7. A computer program product for populating a database with manipulated biological information, said computer program product comprising:
- code for providing a plurality of cells in various stages of the cell cycle, said stages of the cell cycle including at least one selected from interphase, G0 phase, G1 phase, S phase, G2 phase, M phase, prophase, prometaphase, metaphase, anaphase, and telophase;
 - code for manipulating said cells in said various stages of cell cycle development to form a plurality of manipulated cells;
 - code for capturing an image of said plurality of manipulated cells;
 - code for determining a descriptor from said image for said manipulated cells;
 - code for populating a database with said descriptor;
 - wherein said image includes a first component of a cell and a second component of said cell; and
 - a computer readable storage medium for holding the codes.
8. The computer program product of claim 7 wherein said first component and said second component are selected from a protein, a protein modification, a nucleic acid, a lipid, a carbohydrate, a sub-cellular structure and an organelle.
9. The computer program product of claim 7 wherein said image is a digitized representation of said plurality of manipulated cells.
11. The computer program product of claim 9 wherein said digitized representation provides a density value of said plurality of manipulated cells.
11. The computer program product of claim 7 wherein said descriptors comprise numeric or logical values.
12. The computer program product of claim 11 wherein said values further comprises a nucleotide.
13. The computer program product of claim 11 wherein said values further comprises an amino acid letter.
14. A system for capturing images of cells or cell structures, the system comprising:
- a cell holder comprising a plurality of sites in a spatial orientation, each of the sites being capable of holding a plurality of cells to be imaged;

an image capturing device coupled to the cell holder, the image capture device being adapted to capture at least one image in at least one of the plurality of sites;

an illumination apparatus comprising a liquid light guide coupled to the plate for highlighting the plurality of cells in a relatively even spatial manner for image capturing purposes;

an image processing device coupled to the image capturing device, the image capturing device being adapted to convert the image into a digital representation; and

a database storage device comprising a database management element coupled to the image capturing device, the database storage device being adapted to retrieve the digital representation of the image from the image processing device and storing the digital representation.

15. The system of claim 14 further comprising a stage comprising a device for moving the cell holder in a spatial direction to traverse across the cell holder in the spatial orientation.

16. The system of claim 14 wherein the illumination apparatus comprises sub-elements, at least one of the sub-elements being positioned away from the image capturing device to prevent a possibility of vibration from the one sub-elements to be transmitted to the image capturing device.

17. The system of claim 14 wherein the digital representation comprises a plurality of regions and objects.

18. The system of claim 14 further comprising a computing device connected between the database storage device and the image processing device.

19. The system of claim 14 wherein the image capturing device comprises a magnification of at least 1X and greater to capture the image of the site.

20. The system of claim 14 wherein the plurality of sites comprises at least 96 sites.

21. The system of claim 14 wherein the liquid light guide characterized as a flexible member that substantially prevents vibration from the an element of the illumination apparatus to be transferred to the image capturing device.

22. The system of claim 14 wherein the spatial direction can be selected from an x-direction, a y-direction, or a z-direction in a Cartesian coordinate system.

23. The system of claim 14 wherein the each of the sites comprises
5 a volume that is sufficient to prevent a solution therein from evaporating in a substantial manner that may influence the image capturing.

24. A method for identifying a mechanism of action for a first compound, the method comprising the steps of:

receiving the first compound;
10 measuring at least one feature of a cellular phenotype to define a target phenotype;
identifying additional compounds providing a feature similar to the feature identified in the measuring step; and
characterizing the first compound in terms of distance from a specific
15 target phenotype having known characteristics.

25. The method of claim 24 comprising the further step of storing the additional compounds and their associated phenotypes in a database.

26. A method for identifying a mechanism of action for a cellular stimulus, the method comprising the steps of:

20 receiving cells of interest;
measuring at least one feature of the cells to define and quantify a target phenotype;
identifying additional compounds providing a feature similar to the feature identified in the measuring step; and
25 characterizing the first compound in terms of distance from a specific target phenotype having known characteristics.

27. A method for identifying information relevant to at least one of a mechanism of action and cellular activity by utilizing assay data to elucidate a phenotype, the method comprising the steps of:

30 identifying a target protein;
identifying positive and negative biochemical hits related to the target protein;
defining the target phenotype utilizing the positive and negative hits;

identifying other compounds providing similar features; and
characterizing the first compound in terms of distance from a specific
target phenotype having known characteristics.

5




FIG. 1




FIG. 1A




FIG. 1B




FIG. 2




Fig. 3




FIG. 4





FIG 5






FIG. 5B




FIG. 6




Fig. 7A




Fig. 7B
Step 704 of Fig. 7A




Fig. 7C
Step 714 of Fig. 7B




Fig. 7D
Step 716 of Fig. 7B




FIG. 7E




Fig. 7F




Fig 7G




Fig. 7 H




Fig. 71




Fig. 7J




Fig. 7K




Fig. 8A




Fig. 8B




Fig. 8C




Fig. 8D




Fig. 8E





Fig. 8F








Fig. 10








Fig 13




Fig 14




Fig 15




Fig 16




Fig 17

Conversion of morphometric parameters into nucleic acid code
and clustering of the resulting sequences using Neighbor
Joining method.

Compound:	Measurements																							
	Count	Area	Perimeter	Length	Breadth	Fiber length	Fiber breadth	Shape factor	Eli. form factor	Inner radius	Outer radius	Mean radius	Equiv. radius	Equiv. sphere vol.	Equiv. prolate vol.	Equiv. oblate vol.	Equiv. sphere surface area	Average gray value	Total gray value	Optical density	Radial dispersion	Texture Difference Moment	EFA Harmonic 2, Semi-Maj	EFA Harmonic 2, Semi-Min
Control	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	a	t	t	
Taxol	a	t	t	t	t	t	t	t	a	t	t	t	t	t	t	t	t	t	t	t	t	t	t	
CD	c	a	a	a	t	a	t	t	c	a	a	a	a	a	a	a	a	t	a	a	a	t	a	g
Nocodazol	c	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	
Staurosporine	g	g	c	a	a	t	a	a	t	g	a	a	a	t	g	g	g	a	a	t	a	t	a	a
Vinblastine	c	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	g	t	t	t	t	t	
Hydroxyurea	g	t	t	t	t	t	t	g	t	t	t	t	t	t	t	t	t	t	t	c	t	a	t	t




Fig 18

Conversion of morphometric parameters into amino acid codes
and clustering of the resulting sequences using Neighbor
Joining method.

Control	H Count	D Area	T Perimeter	T Length	N Breadth	S Fiber length	O Fiber breadth	W Shape factor	M Ell. form factor	S Inner radius	T Outer radius	T Mean radius	T Equiv. radius	T Equiv. sphere vol.	C Equiv. prolate vol.	C Equiv. oblate vol.	D Equiv. sphere surface a	D Average gray value	M Total gray value	C Optical density	T Radial dispersion	G Texture Difference Mo	T EFA Harmonic 2, Semi-	T EFA Harmonic 2, Semi-	
Taxol	G	F	M	M	P	M	P	H	G	S	M	M	W	C	F	P	F	R	C	M	M	H	M	P	S
CD	F	G	G	G	M	G	M	K	A	G	G	G	G	G	G	G	G	H	G	G	G	M	G	V	H
Nocodazol	W	F	M	M	W	M	P	T	R	S	M	M	M	F	M	W	F	M	M	R	M	M	M	F	G
Staurosporine	N	V	A	G	G	M	G	G	Y	V	G	G	G	M	V	V	V	G	G	H	G	M	G	G	V
Vinblastine	F	W	W	M	W	W	C	W	D	S	M	W	W	M	M	M	W	M	V	E	M	M	M	F	P
Hydroxyurea	S	H	H	H	H	H	H	V	H	H	H	H	H	H	H	H	H	H	H	A	H	G	H	H	D





Fig 19






Fig 21




Fig 22

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
23 November 2000 (23.11.2000)


PCT

(10) International Publication Number
WO 00/70528 A3

- (51) International Patent Classification⁷: G06F 19/00 (72) Inventors: SABRY, James, H.; 52 Buena Vista Terrace, San Francisco, CA 94117 (US). ADAMS, Cynthia, L.; 615 Georgia Avenue, Palo Alto, CA 94306 (US). VAISBERG, Eugeni, A.; 647 Pegasus Lane, Foster City, CA 94404 (US). CROMPTON, Anne, M.; 2 Bellair Place, San Francisco, CA 94133 (US). BLUM, Robert, I.; 17 Shoreview Avenue, San Francisco, CA 94121 (US). OESTREICHER, Donald, R.; 904 Old Town Court, Cupertino, CA 95014-4024 (US). SIGAL, Nolan, II.; 941 Berry Avenue, Los Altos, CA 94024 (US).
- (21) International Application Number: PCT/US00/13154
- (22) International Filing Date: 12 May 2000 (12.05.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- | | | |
|---------------|----------------------------|----|
| 09/310,879 | 14 May 1999 (14.05.1999) | US |
| 09/311,890 | 14 May 1999 (14.05.1999) | US |
| 09/311,996 | 14 May 1999 (14.05.1999) | US |
| 60/134,104 | 14 May 1999 (14.05.1999) | US |
| Not furnished | 26 April 2000 (26.04.2000) | US |
- (74) Agent: LOUIE, Michael, L.; Beyer Weaver & Thomas, LLP, P.O. Box 130, Mountain View, CA 94042-0130 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (71) Applicant: CYTOKINETICS, INC. [US/US]; Suite 2, 280 East Grand Avenue, South San Francisco, CA 94080 (US).

[Continued on next page]

(54) Title: METHOD AND APPARATUS FOR PREDICTIVE CELLULAR BIOINFORMATICS



(57) Abstract: Techniques for using information technology in therapeutics or drug discovery. In an exemplary embodiment, techniques for determining information about the properties of substances based upon information about structure of living or non-living cells exposed to substances are provided. A method according to the present invention enables researchers and/or scientists to identify promising candidates in the search for new and better medicines or treatments using, for example, a cellular informatics database. The present invention further teaches a system for acquiring knowledge from cellular information. The system has a database 1012 comprising a database management module ("DBMS"). The system also has a variety of modules, including a population module coupled to the DBMS for categorizing and storing a plurality of features (e.g., cell size, distance between cells, cell population, cell type) from an image acquisition device into the database. The system has a translation module coupled to the DBMS for defining a descriptor from a set of selected features from the plurality of features. In a specific embodiment, the descriptor is for a known or unknown compound, e.g., drug. A prediction module is coupled to the DBMS for selecting one of a plurality of descriptors from known and unknown compounds from the database based upon a selected descriptor from a selected compound. The selected compound may be one that is useful for treatment of human beings or the like.

WO 00/70528 A3



(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(88) **Date of publication of the international search report:**
8 March 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *With international search report.*

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/13154

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 G06F19/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 G06F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 38490 A (BIODX INC ;DUNLAY R TERRY (US); GOUGH ALBERT H (US); GIULIANO KENN) 3 September 1998 (1998-09-03) cited in the application	1-6, 24-27
Y	page 1; claims 1-43	7-23
X	WO 98 45704 A (TULLIN SOEREN ;KASPER ALMHOLT (DK); NOVONORDISK AS (DK); SCUDDER K) 15 October 1998 (1998-10-15) abstract; claims 1-3,22,73,80,81,86	1-6, 24-27
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *S* document member of the same patent family

Date of the actual completion of the international search

17 November 2000

Date of mailing of the international search report

24/11/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Fillooy García, E

INTERNATIONAL SEARCH REPORT

Inter. onal Application No

PCT/US 00/13154

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MONTIRONI R ET AL: "COMPUTED CELL CYCLE AND DNA HISTOGRAM ANALYSES IN IMAGE CYTOMETRY IN BREAST CANCER" JOURNAL OF CLINICAL PATHOLOGY,GB,LONDON, vol. 46, no. 9, 1 September 1993 (1993-09-01), pages 795-800, XP000644549 ISSN: 0021-9746 abstract	7-13
Y	WO 97 40055 A (DOW CHEMICAL CO ;UNIV TEXAS TECH (US)) 30 October 1997 (1997-10-30) page 18, line 26 - line 32	14-23
P,X	WO 99 39184 A (HARTMANN THOMAS ;RIBOZYME PHARM INC (US)) 5 August 1999 (1999-08-05) the whole document	1-6, 24-27
P,X	WO 00 17643 A (CELLOMICS INC ;DUNLAY R TERRY (US); GOUGH ALBERT H (US); RUBIN RIC) 30 March 2000 (2000-03-30) the whole document	1-6, 24-27
E	WO 00 50872 A (CELLOMICS INC ;KAPUR RAVI (US); GIULIANO KENNETH A (US)) 31 August 2000 (2000-08-31) the whole document	1-6, 24-27
A	GIULIANO K A ET AL: "Fluorescent-protein biosensors: new tools for drug discovery" TRENDS IN BIOTECHNOLOGY,GB,ELSEVIER PUBLICATIONS, CAMBRIDGE, vol. 16, no. 3, 1 March 1998 (1998-03-01), pages 135-140, XP004108592 ISSN: 0167-7799 page 139, left-hand column, paragraph 4 -right-hand column, paragraph 3	1-27

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/13154

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9838490 A	03-09-1998	US 5989835 A	23-11-1999
		US 6103479 A	15-08-2000
		AU 6667898 A	18-09-1998
		EP 0983498 A	08-03-2000
		JP 2000509827 T	02-08-2000
		AU 3297197 A	05-01-1998
		EP 0912892 A	06-05-1999
WO 9845704 A	15-10-1998	AU 6820998 A	30-10-1998
		EP 0986753 A	22-03-2000
WO 9740055 A	30-10-1997	AU 719002 B	04-05-2000
		AU 2660697 A	12-11-1997
		CA 2251924 A	30-10-1997
		EP 0898574 A	03-03-1999
		NO 984840 A	16-12-1998
		US 5928627 A	27-07-1999
WO 9939184 A	05-08-1999	AU 2241899 A	16-08-1999
WO 0017643 A	30-03-2000	AU 6048599 A	10-04-2000
WO 0050872 A	31-08-2000	NONE	